

Mapping Allergenic Pollen Vegetation in UK to Study Environmental Exposure and Human Health

Rachel N. McInnes^{a,b,*}, Deborah Hemming^{a,c}, Peter Burgess^d, Donna Lyndsay^e, Nicholas J. Osborne^{f,b}, Carsten Ambelas Skjøth^g, Sam Thomas^h, Sotiris Vardoulakis^{i,b}

^aMet Office Hadley Centre, FitzRoy Road, Exeter, EX1 3PB, UK

^bEuropean Centre for Environment and Human Health, University of Exeter Medical School, Knowledge Spa, Royal Cornwall Hospital, Truro, TR1 3HD, UK

^cBirmingham Institute of Forest Research, School of Geography, Earth and Environmental Sciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK

^dDevon Wildlife Trust, Cricklepit Mill, Commercial Road, Exeter, EX2 4AB, UK

^eBluesky International Limited, Unit 3, Jackson Street, Coalville, Leicestershire, LE67 3NR, UK

^fSchool of Public Health and Community Medicine, University of New South Wales, Sydney, New South Wales, 2052, Australia

^gNational Pollen and Aerobiological Research Unit, Institute of Science and the Environment, University of Worcester, WR2 6AJ, UK

^hInstitute of Biological, Environmental & Rural Sciences (IBERS), Aberystwyth University, Penglais, Aberystwyth, Ceredigion, SY23 3DA, UK

ⁱEnvironmental Change Department, Centre for Radiation, Chemical and Environmental Hazards, Public Health England, Chilton, Oxon, OX11 0RQ, UK

Abstract

Allergenic pollen is produced by the flowers of a number of trees, grasses and weeds found throughout the UK. Exposure to such pollen grains can exacerbate pollen-related asthma and allergenic conditions such as allergic rhinitis (hay fever). Maps showing the location of these allergenic taxa have many applications: they can be used to provide advice on risk assessments; combined with health data to inform research on health impacts such as respiratory hospital admissions; combined with weather data to improve pollen forecasting systems; or as inputs to pollen emission models. In this study we present 1 km resolution maps of 12 taxa of trees, grass and weeds found in the UK. We have selected the main species recorded by the UK pollen network. The taxa mapped in this study were: *Alnus* (alder), *Fraxinus* (ash), *Betula* (birch), *Corylus* (hazel), *Quercus* (oak), *Pinus* (pine) and *Salix* (willow), *Poaceae* (grass), *Artemisia* (mugwort), *Plantago* (plantain), *Rumex* (dock, sorrels) and *Urtica* (nettle). We also focus on one high population centre and present maps showing local level detail around the city of London. Our results show the different geographical distributions of the 12 taxa of trees, weeds and grass, which can be used to study plants in the UK associated with allergy and allergic asthma. These maps have been produced in order to study environmental exposure and human health, although there are many possible applications. This novel method not only provides maps of many different plant types, but also at high resolution across regions of the UK, and uniquely present 12 key plant taxa using a consistent methodology. To consider the impact on human health due to exposure of the pollen grains, it is important to consider the timing of pollen release, and its dispersal, as well as the effect on air quality, which is also discussed here.

Keywords: Aeroallergen, Allergenic pollen, Human health, Land cover, Source map, Species distribution

*Corresponding author: rachel.mcinnes@metoffice.gov.uk

1. Introduction

Allergenic pollen is produced by a number of trees, grasses and weeds found throughout the United Kingdom (UK). Exposure to such pollen grains can result in exacerbation of pollen-related asthma and allergenic conditions such as allergic rhinitis (pollenosis or hay fever)(Greiner et al., 2012). With the total effect of future environmental changes on pollen production and spread being highly uncertain (Osborne and Eggen, 2014), there is a need for detailed and robust pollen-related health impact information. Highly detailed maps of locations of allergenic plants in the UK, as presented here, are an important part of this as they provide the spatial detail required for impact assessments. Such maps will be highly useful for the next generation co-exposure modelling system currently under development for the UK area. This system is based on an extension of WRF-Chem model (e.g. Grell et al. (2011)) and is designed for handling both chemical air pollutants (e.g. Werner et al. (2016, 2017)) and bioaerosols (e.g. pollen and spores) at the species level (Skj oth et al., 2015a) where both chemical air pollutants and bioaerosols directly interact with and feedbacks to atmospheric physics.

The UK has one of the highest prevalence of doctor diagnosed asthma affecting around 10% of the adult population (Netuveli et al., 2005). Approximately 80% of people with asthma also have a pollen allergy (Asthma UK, 2017). The NHS Choice website states that over 10 million people have hay fever in England (NHS Choices, 2017). Physician-based diagnosis of allergic rhinitis 13.2% (95% CI 11.6-14.9) in the UK in 2001 (Bauchau and Durham, 2004).

Detailed maps of allergenic pollen producing taxa, in combination with pollen forecasts and calendars, can help sufferers to manage their condition by reducing their exposure. Asthma and allergic rhinitis significantly reduce quality of life and have a large economic impact (Bousquet et al., 2001). As such, it is a significant environmental health issue.

Our method is novel as it not only provides maps of many different plant types, but also at high resolution across regions of the UK. These maps fill a need for detailed vegetation mapping of allergenic plant species across a whole country, to improve understanding of relationships between allergenic pollen exposure and human health outcomes. With an initial assessment of data availability and expertise on plant types and land cover, the method presented here could be duplicated for other locations around the world.

In this paper we present detailed maps of location of different taxa monitored by the UK pollen network that are associated with allergy and allergic asthma. These have been made to study environmental exposure and human health, although these new vegetation maps have many possible applications as outlined in the following section.

1.1. Purpose of this work and wider context

Vegetation maps have many applications worldwide: they can be used to provide advice on risk assessments, e.g. on invasive species (for example Csornai et al. (2011)); combined with health data to inform research on health impacts such as respiratory hospital admissions caused by exposure of environmental aeroallergens (Newson et al., 2014; Bousquet et al., 2007); combined with weather data to improve pollen forecasting systems (e.g. Zink et al. (2012)); or as inputs to dedicated pollen emission models (Zink et al., 2013).

69 Vegetation maps have many applications worldwide: they can be used to provide advice on risk assessments;
70 combined with health data to inform research on health impacts such as respiratory hospital admissions;

71 The vegetation maps presented here have been developed to study human exposure to allergenic pollen, and the
72 effect on human health. They can be combined with hospital admissions data for asthma to study the impact of
73 different allergenic taxa in the UK. By producing detailed maps of the location of allergenic pollen producing plants
74 for the UK it may be possible to identify the taxa that increase risk of higher hospital admissions for asthma in that
75 particular region. This level of detail could help with the accurate measurement of health impacts as well as monitoring
76 for climate impacts through changes in vegetation distribution, and changes to pollen allergenicity.

77 Vegetation mapping of plants with allergenic pollen may also help affected individuals with self-management of
78 their allergy or asthma. Once linked with health effects, another application of these maps will be to provide advice
79 for vegetation management practices. These practices can include the choice of tree species, sex of tree for planting,
80 and grass cutting regimes to limit exposure to the most allergenic pollen.

81 Detailed maps of the location of pollen producing plants are also key to a future pollen forecasting system. Maps
82 can be coupled with key weather variables such as wind direction and speed, precipitation, humidity and temperature,
83 and both phenological and dispersion models, to predict the emission timing, amount (Skjøth et al., 2012) and disper-
84 sion of pollen grains, to provide more spatially and temporally resolved pollen forecasts [for the UK, compared to the](#)
85 [existing regression based approaches for trees \(Adams-Groom et al., 2002\), grasses \(e.g. Smith and Emberlin \(2005,](#)
86 [2006\)\) and weeds.](#)

87 *1.2. Health impacts of allergenic pollen*

88 Exposure to allergenic pollen from certain trees, grasses and weeds is associated with a range of health effects, in-
89 cluding allergic rhinitis (hay fever), exacerbation of asthma in susceptible individuals, and atopic dermatitis (eczema)
90 (Cecchi, 2012). These pollen grains are produced in the flowers of angiosperm plants (e.g. most deciduous trees,
91 weeds and grasses), and in the pollen cones of gymnosperm plants (e.g. conifer trees), and the timing of their re-
92 lease varies depending on the species and environmental conditions. Susceptible individuals can be sensitive to pollen
93 from one or more different type of trees, grasses or weeds. [Estimates of the levels of allergies towards environmental](#)
94 [aeroallergens \(pollen, spores and cat/dog/house dust mites\) among patients range typically from up to 30%-50% in](#)
95 [Europe \(Newson et al., 2014\), where the largest fractions of sensitisations towards specific pollen typically are against](#)
96 [grasses and trees of the *Betulaceae* family with clinically relevant sensitisation rates for grasses exceeding 50% for](#)
97 [both UK and Denmark \(e.g. Burbach et al. \(2009\)\).](#)

98 *1.3. Background to allergenic vegetation mapping*

99 Here we outline methods used in the literature to produce spatial maps detailing the location of vegetation that
100 emits allergenic pollen, often referred to as pollen source maps, or inventories (Skjøth et al., 2012). ‘Bottom up’
101 and ‘top down’ approaches to creating pollen source location maps are outlined in Skjøth et al. (2012). ‘Bottom up’

102 techniques start with the location and amount of the pollutant — they combine land cover maps with regional scale
103 statistics, for example as used in Skjøth et al. (2008). Availability of these regional scale statistics is what limits
104 this approach, for example statistical information on the regional abundance of weeds and certain taxa is largely not
105 available. Many sources exist which refer to the presence or absence of a particular plant type or species, but not
106 their abundance, and these rarely cover a whole area, for example the size of the UK. ‘Top down’ inventories start
107 with a measured quantity and work backwards using models to calculate the distribution. They use station based
108 pollen monitoring observations along with land cover maps, for example Skjøth et al. (2012) which uses data from
109 the European Aeroallergen Network (EAN). Geographic and temporal coverage of monitoring stations are however
110 limiting factors of ‘top down’ methodologies. Existing pollen mapping includes tree source maps at 50 km × 50 km
111 resolution (Skjøth et al., 2008) and *Ambrosia* (ragweed) source inventories for different European locations (Skjøth
112 et al., 2010; Thibaudon et al., 2014; Karrer et al., 2015). Statistical mapping of trees over Europe at 1 km × 1 km
113 resolution is presented in Brus et al. (2011). Skjøth et al. (2015b) used back-trajectories of pollen observations to
114 create footprint areas of *Alnus* (alder) and *Betula* (birch) in Worcester. Remote sensing can also be used such as
115 Skjøth et al. (2013) who created a grass pollen inventory in Aarhus (Denmark) using GIS and remote sensing.

116 The quality of the mapping is dependent on resolution and taxa specificity of available input datasets. High
117 resolution inputs can provide a challenge for handling large volumes of data, and lower resolution inputs may be
118 easier to obtain, but can miss important spatial detail.

119 In this paper we present a novel set of maps of key plants associated with allergy and allergic asthma in the UK.
120 They are provided at higher resolution coverage than currently available elsewhere in the literature, use a state-of-the-
121 art tree dataset (Bluesky, National Tree Map, 2015) for highly accurate tree locations, and uniquely present 12 key
122 plant taxa (monitored by the UK pollen network), using a consistent methodology.

123 **2. Materials and methods**

124 *2.1. Selection of allergenic vegetation*

125 Tree and weed taxa and grasses were selected which were monitored by the UK pollen network (run by the Met
126 Office (Met Office, 2017b)), where the majority are considered medium to highly allergenic, and for which datasets
127 could be produced covering Great Britain. As these include either species i.e. *Acer pseudoplatanus* (sycamore) or
128 genera i.e. *Alnus* (alder), for clarity in the following text we will refer to all as ‘taxa’.

129 A summary of the availability of tree taxa information present in the Forestry Commission’s National Forest
130 Inventory (Forestry Commission, Great Britain, 2011), and the Trees in Towns II report (Britt and Johnston, 2008) —
131 both of which were used in this study is given in Table 1. The seven tree taxa which had data availability that were
132 selected to be studied in this paper, are indicated in bold font in Table 1. Note that the seven selected tree taxa include
133 those which were reported to be most allergenic by the review by de Weger et al. (2012) — *Corylus* (hazel), *Alnus*

134 (alder), *Betula* (birch) and *Quercus* (oak) — excluding *Platanus* (plane) and those allergenic but not prevalent in the
135 UK, *Olea* (olive) and *Cupressaceae* (cypress).

136 All grasses, *Poaceae*, were considered together as the current UK pollen monitoring observations are unable to
137 delineate between the various species, also very little is known about which species of grass are most allergenic in
138 the UK. Note that, in Europe, grass pollen is the most important pollen allergen due to both its large distribution in
139 Europe (Skjøth et al., 2012) and its record high sensitisation rates among patients (Burbach et al., 2009).

140 The final group of taxa looked at fall into the category of ‘weeds’. The National Pollen and Aerobiological
141 Research Unit (University of Worcester) identify five plants in this category, all of which are monitored in the UK
142 as part of the pollen network: *Rumex* (dock, sorrels), *Artemisia* (mugwort), *Urtica* (nettle), *Plantago* (plantain) and
143 *Brassica napus* (oil seed rape). Oil seed rape is planted in the UK for agriculture, and due to a lack of detailed mapping
144 information on the spatial abundance of it in the UK, we omitted this plant from this study.

145 Summarising, the 12 selected taxa from the UK pollen network mapped in this study were as follows. **Trees:**
146 *Alnus* (alder), *Fraxinus* (ash), *Betula* (birch), *Corylus* (hazel), *Quercus* (oak), *Pinus* (pine) and *Salix* (willow). **Grass:**
147 *Poaceae* (bulk grass, not species specific). **Weeds:** *Artemisia* (mugwort), *Plantago* (plantain), *Rumex* (dock, sorrels)
148 and *Urtica* (nettle).

149 2.2. Grass mapping methodology

150 The Centre for Ecology and Hydrology (CEH) Land Cover Map 2007 (hereafter LCM 2007) is a 25 m resolution
151 raster map of Great Britain which shows land cover using 23 different classifications (Morton et al., 2011). This was
152 used to create the grass map in this study. Land cover classes where grasses are predominantly found were selected.
153 The six grassland classes used are shown in Table 2. Note that some other categories — namely, Arable, Urban and
154 Suburban (classes 3, 22 and 23 respectively) are likely to contain some grass, but due to a lack of detailed spatial
155 information these areas have not been included. These were not selected for this study. See Centre for Ecology &
156 Hydrology, UK (2011) for full description of the land cover classes and their habitats. Raster masks of the selected
157 categories were made and applied using a script written in the programming language PYTHON (Python Software
158 Foundation, 2017). This resulted in a 25 m raster map showing only grassland areas. The grass map was then
159 regrided to 1 km resolution and converted to percentage cover, using the spatial analyst toolbox in software ARCGIS
160 DESKTOP VERSION 10 (ESRI, 2017).

161 2.3. Weed mapping methodology

162 Statistical information on the regional abundance of allergenic weeds is not consistently available across the UK.
163 Hence, we have developed a method based on that used by Skjøth et al. (2010); Kaye and Burgess (2013); Thibaudon
164 et al. (2014) and Karrer et al. (2015) which uses land cover map classes and input of expert plant knowledge to filter
165 for habitat selection. Here we used expert elicitation to determine the suitability of a habitat (land cover class) for each

tree genera or species — taxa	common name	broadleaf or conifer	pollen network	NFI	Trees In Towns II
<i>Acer pseudoplatanus</i>	sycamore	broadleaf	—	✓	✓
<i>Alnus</i>	alder	broadleaf	✓	✓	✓
<i>Betula</i>	birch	broadleaf	✓	✓	✓
<i>Castanea</i>	chestnut	broadleaf	—	✓	—
<i>Corylus avellana</i>	common hazel	broadleaf	✓	✓	✓
<i>Crataegus</i>	hawthorn	broadleaf	—	✓	✓
<i>Eucalyptus</i>	gum	broadleaf	—	—	✓
<i>Fagus</i>	beech	broadleaf	—	✓	✓
<i>Fraxinus</i>	ash	broadleaf	✓	✓	✓
<i>Larix</i>	larch	conifer	—	✓	—
<i>Picea</i> *	spruce	conifer	—	✓	✓
<i>Pinus</i>*	pine	conifer	✓	✓	✓
<i>Platanus</i>	plane	broadleaf	✓	—	✓
<i>Populus</i>	poplar	broadleaf	✓	—	✓
<i>Pseudotsuga</i>	douglas fir	conifer	—	✓	—
<i>Quercus</i>	oak	broadleaf	✓	✓	✓
<i>Salix</i>	willow	broadleaf	✓	✓	✓
<i>Taxus</i>	yew	conifer	✓	—	✓
<i>Tilia</i>	lime	broadleaf	✓	—	✓
<i>Ulmus</i>	elm	broadleaf	✓	—	✓

Table 1: Summary of the availability of data for tree genera or species — collectively referred to in the text as taxa — from the Forestry Commission National Forest Inventory (NFI) report (Forestry Commission, Great Britain, 2011) and the Trees in Towns II report (Britt and Johnston, 2008). The tree taxa mapped in this study were chosen as those which were monitored as part of the UK pollen network and appeared in both the NFI and Trees in Towns II data, highlighted in the table in bold font. ‘*’ denotes where specific species are recorded: *Picea* (spruce) — *sitchensis* (sitka spruce), *abies* (Norway spruce); *Pinus* (pine) — *sylvestris* (Scots pine), *nigra* (Corsican pine), *contorta* (lodgepole pine).

LCM 2007 class	class number
improved grassland	4
rough grassland	5
neutral grassland	6
calcareous grassland	7
acid grassland	8
heather grassland	11

Table 2: LCM 2007 classes selected as grassland, along with their classification number in the LCM 2007 dataset.

weed. The type of expert elicitation used was ‘behavioural aggregation’ — group decision making — where experts pooled knowledge (O’Hagan et al., 2006) to reach a consensus.

Three experts with documented specialist knowledge on the selected weeds were chosen and each provided their individual independent assessment of ‘habitat suitability’ and percentage cover of the weeds in each land class category of the LCM 2007 dataset. For each taxon, each expert scored each land class 0,1 or 2 — 0 indicated that the habitat was not suitable for this taxon, 1 that the habitat was mildly suitable, and 2 that it was a largely suitable habitat. Using this ‘habitat suitability’ indicator as a guide, the experts then individually and independently assigned an estimate of ‘fractional cover’ for that taxon in that habitat. For example, for a habitat suitability of ‘2’ (largely suitable habitat) of *Plantago* in land class ‘urban’, each expert estimated a typical area coverage in that land area of *Plantago*. Following a similar approach to the ‘Delphi method’ (O’Hagan et al., 2006), each expert first provided their own independent assessment. This was then supplied to all the other experts in the group, and the experts were given the chance to update their assessment. The experts were also asked to make any relevant notes regarding land classes and these taxa, in order to capture any key aspects to take into account with the mapping. After having the independent assessments from each expert, the experts then collaboratively reached a final value for the percentage cover estimates (‘behavioural aggregation’). The experts collectively discussed each weed taxon and came to a joint agreement on value for each percentage cover in each of the land classes. Values of percentage cover for typical 25 m × 25 m grid box of each land class ranged between: mugwort 0-0.1%; plantain 0-5.0%; *Rumex* 0-2.0%; nettle 0-1.0%.

A 25 m percentage cover map was created for each of the four weeds, by using ‘reclassify’ tool in ArcGIS to apply taxa coverage to different land classifications. The 25 m map was then aggregated to 1 km using the same technique used for grass, as described above.

Note that there are differing numbers of taxon in each genus and that each of these will vary in terms of their relative abundance by land class. This will not overly affect the outcomes as most of the species are either rare or not present in the UK. For example, there are 57 species of *Artemisia* in the north temperate regions (Grewling et al., 2012), but only *A. vulgaris* (mugwort), *A. campestris* (field wormwood) and *A. absinthium* (wormwood) are

190 widespread throughout Europe. The online Atlas of the British and Irish flora lists 10 different *Artemisia* species, but
191 only wormwood and mugwort have widespread records. A consequence is that the vast majority of pollen related to a
192 specific genera will originate from a few common species.

193 2.4. Tree mapping methodology

194 A high resolution tree map was used to build a base map of all tree locations. Scaling factors were created which
195 accounted for the proportion of broadleaf and conifer trees at different locations. Finally, regional taxa fractions were
196 applied. The final taxa maps were created by combining all of these data. This combination can be described in the
197 following way to calculate the number of trees of a given taxon in a grid box:

$$T_{\text{taxon}} = \alpha_{\text{taxon}} \times \beta \times T_{\text{all}}, \quad (1)$$

198 where T_{all} is the total number of all trees in a grid box, α_{taxon} is the fraction of that taxon found in that grid box
199 (a function of region/urban area), and β is the scaling factor for broadleaf/conifer abundance in a given land type (a
200 function of region and also different for broadleaf/conifer taxa).

201 The method used to assign each of these terms is described in detail below.

202 2.4.1. Base map, T_{all}

203 The Bluesky International, National Tree Map (Bluesky, National Tree Map (2015), hereafter NTM) is a dataset
204 built from high resolution aerial photography and remote sensing, and contains the location, height and canopy/crown
205 extents for every single tree which is over 3 m in height. NTM does not contain any species-level or taxon information
206 for the trees. It covers England and Wales in extent. The dataset is provided as a vector GIS dataset, with height and
207 area attributes for each tree. The number of trees (count) in 25 m \times 25 m grid cells was calculated from the NTM
208 dataset using a PYTHON script, with the grid aligned to that of the Forestry Commission dataset. This map formed a
209 ‘base map’ which showed the locations of all trees (over 3 m), to which taxon fractions would be applied. This base
210 map provides the value of T_{all} in Eq. 1, the total number of all trees in a given grid box.

211 2.4.2. Taxon fractions, α_{taxon}

212 Regional measures of taxa totals in Great Britain are presented by the Forestry Commission for broadleaved trees
213 (Brewer, 2013) and coniferous trees (Forestry Commission, Great Britain, 2012). To calculate taxa fractions in forests
214 for the broadleaved taxa, data from *Number of trees by principal broadleaved species for National Forest Inventory*
215 *regions* Brewer (2013, Table A.7) were used to calculate the proportion of each taxon as a fraction of total number of
216 ‘all broadleaves’. A measure of the number of trees per taxon per region was not available for coniferous taxa, and
217 so taxon fractions for pine were instead calculated using the timber volume measure. Data from *Standing coniferous*
218 *timber volume by principal species for National Forest Inventory regions* Forestry Commission, Great Britain (2012,
219 Table 9) were used to calculate the proportion of pine as fraction of total volume of ‘all conifers’ in forests. In urban

tree type	location	taxa fraction source	spatial variation
broadleaf	forests	Forestry Commission 2013	regional
conifer	forests	Forestry Commission 2012	regional
all	urban	Trees in Towns II	none
broadleaf	elsewhere	Forestry Commission 2013	regional
conifer	elsewhere	Forestry Commission 2012	regional

Table 3: Source of taxa fraction data for broadleaf and conifer trees in different land types (forests, urban, elsewhere). The final column outlines the spatial variation of these taxa data. ‘Forestry Commission 2013’ denotes data from Brewer (2013, Table A.7) — fractions were calculated per taxon as fraction of number of ‘all broadleaves’. ‘Forestry Commission 2012’ denotes data from Forestry Commission, Great Britain (2012, Table 9) — fractions were calculated per taxon, as a fraction of total volume of ‘all conifers’. In urban areas, tree taxa fractions were calculated using Trees in Towns II report Britt and Johnston (2008, Figure 2.18). N.B. *Corylus* (hazel) was not included in Figure 2.18, and so a taxa fraction for hazel was obtained from Britt and Johnston (2008, Table 2.8). Vegetation type fractions calculated in urban areas are given as a fraction of total number of *all trees*.

220 areas, tree taxa fractions were calculated using Trees in Towns II report Britt and Johnston (2008, Figure 2.18), a
221 large study of urban trees carried out in towns with populations over 3000. The distribution of taxa in towns differed
222 sufficiently from the forest data that we used these taxa fractions in urban areas. Note that *Corylus* (hazel) was not
223 included in this Figure 2.18, and so a fraction for hazel was obtained from Britt and Johnston (2008, Table 2.8), other
224 fractions calculated from Figure 2.18. The remaining areas — those which were neither forest, nor towns — were
225 grouped as the ‘elsewhere’ category, and in these areas the Forestry Commission taxa fractions, as calculated for
226 forests were used. We considered Countryside Survey data (Maskell et al., 2013) which contained some data on some
227 of the key allergenic taxa in non-forested countryside areas. However, it did not have sufficient information for us to
228 calculate taxa fractions¹. Table 3 presents a summary of the sources of taxa fraction data for broadleaf and conifer
229 trees in different land types (forests, urban, elsewhere). Also shown is spatial variation (final column) of these taxa
230 data — which is regional for all but the urban data.

231 The values of taxon fraction described above provide the value of α_{taxon} in Eq. 1, the fraction of a given taxon in
232 a grid box.

233 2.4.3. Scaling factor, β

234 A ‘scaling factor’ was necessary to account for the fact that many of the taxa fractions were available as a pro-
235 portion of either all broadleaves or all coniferous taxa, and not as a proportion of total number of trees. Differing

¹Although the Countryside Survey had estimates of areas covered by the above taxa, and upper and lower confidence limits on those, it does not provide, as the Trees in Towns did, either an ‘everything else’ category for trees beyond the ‘principal species’, nor does it give the total amount of coverage of all trees, so it is not possible to calculate accurate taxa fractions.

236 proportions of broadleaf and conifer trees are found in different land types and forest types, and so these classifi-
237 cations were used to scale the tree map in different areas, depending on whether the tree type being mapped was
238 classed as broadleaved (alder, ash, birch, hazel, oak and willow) or coniferous (pine). The scaling for broadleaved or
239 coniferous trees in different land types is outlined below.

240 Forest boundaries from Forestry Commission National Forest Inventory (NFI) map (25 m × 25 m resolution;
241 Forestry Commission, Great Britain (2011)) were used to define forested areas. Four NFI interpreted forest types
242 were used: broadleaf; mixed mainly broadleaf; mixed mainly conifer; conifer. Table 4 shows the NFI interpreted
243 forest types, the range of broadleaf and conifer trees expected in each (as defined by Forestry Commission, Great
244 Britain (2011)) and the scale factors, β used for each, for broadleaf and conifer trees.

245 LCM 2007 ‘urban’ and ‘suburban’ classes were used to define urban areas. They were selected to be continuous
246 areas ≥ 50 ha in size, with populations ≥ 3000 (selected using Office of National Statistics (Office of National Statistics,
247 2017) population dataset), not defined as forest in the NFI forestry dataset, and not a thin motorway or airport type
248 infrastructure. Because the Trees In Towns II report used for taxa data included urban parks (for example Regent Park,
249 London) these were included in the urban category here, by including any areas which were completely surrounded
250 by urban/suburban land classes. In Trees In Towns II, each taxa is given as a proportion of all trees, and so no scaling
251 factor is needed (i.e. a multiplicative scaling factor of value 1 was used), as shown in Table 4.

252 The remaining areas — those which were neither forest, nor towns — were grouped as the ‘elsewhere’ category. A
253 calculation of broadleaf and conifer proportions was carried out, using data from Forestry Commission, Great Britain
254 (2011, Table 3). Values of total area (ha) of broadleaf and conifer trees were used and a relative proportion of each
255 calculated from this. Because of the regional differences seen in the forestry data, the highest resolution data available
256 were used for this — in this case England and Wales (rather than GB as a whole). The result was 71% broadleaf and
257 29% conifer in England, and 48% broadleaf and 52% conifer in Wales. These values were used as the scaling factor,
258 β for ‘elsewhere’ areas, as shown in the last two rows of Table 4.

259 2.4.4. *Combining to create final maps*

260 For each of the individual tree taxa, the base map, scaling factor and taxa fractions were combined as illustrated
261 in Figure 1. Note that our tree maps cover England and Wales only because of the data availability of Bluesky NTM.
262 This figure demonstrates the spatial variation in each of the values of the components of Eq. 1 — with the scale factor
263 being a function of land type (i.e. type of forest, and variation in ‘elsewhere’ category in England and Wales) and
264 the proportions of broadleaf/conifers expected there, and the taxa fraction being a function of region and urban areas.
265 The base tree map has a lot of spatial structure, and is in units of number of trees per grid box, the same units as the
266 resulting taxon map. These maps were then regridded to 1 km × 1 km resolution, for comparison with other data and
267 be more manageable in size.

land type	broadleaf cover	conifer cover	scale factor	broadleaf	conifer
<i>forest:</i>					
broadleaf	>80%	<20%	β_b	0.9	0.1
mixed (b)	50%-80%	20%-50%	β_{mb}	0.65	0.35
mixed (c)	20%-50%	50%-80%	β_{mc}	0.35	0.65
conifer	<20%	>80%	β_c	0.1	0.9
urban	-	-	β_u	1	1
elsewhere (E)	71%	29%	β_e	0.71	0.29
elsewhere (W)	48%	52%	β_e	0.48	0.52

Table 4: National Forest Inventory interpreted forest types and range of areas covered by broadleaf and conifer trees in these categories ('broadleaf cover', 'conifer cover'). Note that forest type 'mixed (b)' denotes a forest with mixture of broadleaf and conifer trees, but with majority broadleaf. Similarly 'mixed (c)' contains mixed trees, mainly conifer. Also shown is the scale factor β assigned to each forest/land type, and value of this scale factor for broadleaf and conifer trees (final two columns). The taxa fraction in urban areas is calculated as a proportion of all trees, and so no scaling factor is needed (i.e. a multiplicative scaling factor of value 1 was used). Land types 'elsewhere (E)' and 'elsewhere (W)' denote whether the 'elsewhere' (non-forest, non-urban) area was in England or in Wales respectively, as differing broadleaf/conifer proportions were used.

268 2.5. Alternative datasets and areas

269 In this study we used datasets which are specific to the UK, or parts of it. In order to apply our methodology to
270 different geographical areas, one must identify suitable input datasets. We recommend that one assesses the relevant
271 merits of land use datasets (land classifications) in the desired area of study. Things to consider when selecting a land
272 use dataset would be accuracy, age, resolution and coverage of the dataset, as well as any specific merits / caveats
273 of each dataset which might relate to the specific type of vegetation being studied. Some examples of large land
274 use datasets are Corine (Corine, 2017, 100 m resolution, covering most of Europe;), ESA Landcover map (ESA,
275 2017, 300 m resolution, global coverage), IGBP (2017), and land cover data from satellite observations such as SPOT
276 VEGETATION (Spot Vegetation, 2017) and MODIS Vegetation Continuous Fields (2017). Similarly, one must assess
277 the availability and quality of forestry or tree information, and also that of taxon information provided at a regional
278 (or higher) level. One such dataset which is available over Europe is the European Forest Inventory (EFI) dataset
279 (Schelhaas et al., 2006) which has taxa level maps at 1 km resolution. In this study we considered the EFI but found
280 the EFI total tree map (all taxa) differed significantly from the Bluesky International NTM dataset on tree location,
281 which we believed to be a more accurate indication of tree location due to its methodology. For this reason we used
282 just the NTM dataset in further analysis. Thus caution would be urged if using EFI in place of a higher resolution and




Figure 1: Diagram illustrating method used to create tree taxa maps.

283 more accurate tree dataset (e.g. up-to-date aerial or satellite based dataset with a common methodology across the
 284 dataset), as it will affect the quality of the resulting maps.

285 We note that without access to such a high resolution dataset such as the Bluesky International NTM dataset on
 286 tree location, this will have an impact on the quality of the resulting maps. One must consider changing the resolution
 287 of the final maps with respect to the quality of the input maps, and not provide higher resolution maps than the data
 288 reasonably allow. This judgement must be made by those familiar with land cover and vegetation datasets and their
 289 strengths and weaknesses, and considering what the final maps will be used for.

290 For this reason we have not produced tree taxon maps in Scotland where the Bluesky International NTM maps were
 291 not available. This decision was made because we wanted to use a consistent methodology and produce consistent
 292 output such that the provided taxon maps could be used for human health studies.

293 3. Results

294 Here we present 1 km gridded maps of the 12 selected taxa.

295 3.1. Grass map

296 Figure 2 shows the resulting *Poaceae* (grass) location map (percentage coverage), gridded at 1 km resolution, for
 297 Great Britain. The colourbar shows equal sized quantiles. An overall pattern of higher percentage cover of grass in the
 298 northern and western regions of Great Britain, and lower overall cover to the south east can be seen. High mountainous
 299 regions in Scotland show lower levels of grass cover than their surrounding lower lying areas. The eastern central area
 300 of England has the lowest percentage grass coverage, while Wales and north west England, and western Scotland have
 301 the highest density of grass coverage. Note that these maps show the vegetation location of grass, and not of the pollen
 302 concentration, we discuss how different factors affect the spread of pollen grains in Section 4.




Figure 2: Map showing location of *Poaceae* (grass), in units of percentage cover per 1 km × 1 km grid box, across Great Britain. Colour ranges have been set to 5 equal sized quantiles (i.e. each contains 20% of data). Image Crown Copyright, 2016, The Met Office. Based on digital spatial data licensed from the Centre for Ecology & Hydrology, copyright NERC (CEH).

303 3.2. Weed maps

304 Figure 3 shows the location maps (percentage coverage) for the four selected weed taxa, gridded at 1 km resolution,
305 for Great Britain with quantile based colourbars, allowing a comparison of the relative locations of each taxon across
306 Great Britain. Note that the total quantities are different for each weed taxon. The same figure without quantile-based
307 colourbars are shown in the Supplementary materials, Figure A.8. Dock (*Rumex*, includes dock and sorrels) is fairly
308 evenly distributed across Great Britain, with no overall north/south or east/west trend. It is not abundant in urban areas,
309 or in high altitudes, and these areas can be seen in white and grey on the map. Nettle distribution display a generally
310 lower values of percentage cover than dock, and shows a north/west - south/east trend, with higher abundance in
311 southern and eastern Great Britain, largely a result of lower cover at higher altitude. Unlike dock, nettle has its highest
312 coverage in urban areas. Plantain is relatively evenly distributed across the whole of Great Britain, with highest areas
313 of coverage in urban areas. Mugwort has much lower general levels of coverage, reaching a maximum of only 0.1%.
314 Mugwort has a similar distribution to nettle, being found mostly in the south and eastern areas of Great Britain, and
315 in urban areas.

316 3.3. Tree maps

317 Tree taxon plots for the 7 tree taxa are shown in Figures 4 and 5. Units are in number of trees per 1 km × 1 km grid
318 square. These figures have quantile based colourbars, allowing a comparison of the relative locations of each taxon in
319 England and Wales. Note that the total quantities are different for each tree taxon. The same figures without quantile-
320 based colourbars are shown in the Supplementary materials Figures A.9 and A.10. Of the broadleaf trees, alder and
321 willow have the lowest number of trees, followed by oak and ash. Birch and hazel both have higher abundance in
322 terms of number of trees. Pine is the only conifer tree mapped in this study, and has much higher number of trees
323 than the broadleaves mapped here. In England, pine is the main conifer taxon found, and it is very abundant. The
324 pine map shows a different pattern of distribution to that of the broadleaves, notably a lack of pine in greater London
325 is seen, and less pine trees in Wales and western England. The highest concentrations of hazel are found in the south
326 east of England, and lowest values in the eastern area of England. Birch also has peaks in the south east of England.
327 The distributions of oak and ash are very similar to each other; while both the alder and willow maps show a peak in
328 western areas of England and Wales, as well as the south east of England. The results here are not directly comparable
329 with those in Skjøth et al. (2008), because our maps are much higher resolution, and include all trees — not only those
330 in forests.

331 3.4. Focus on high population centre

332 Maps showing the allergenic plant locations across areas of high population are valuable because the large number
333 of people living in these cities means that many are exposed to pollen from these plants. Figure 6 shows the percentage
334 cover of the four weed taxa, and grass, across greater London. Figure 7 presents the percentage cover of the 7
335 allergenic trees in this same area.




Figure 3: Maps showing location of four allergenic weeds in Great Britain, in units of percentage cover per 1 km × 1 km grid box. ¹⁵ *Top left*: percentage cover of *Rumex* (dock, sorrels); *Top right*: percentage cover of *Urtica* (nettle); *Bottom left*: percentage cover of *Plantago* (plantain); *Bottom right*: percentage cover of *Artemisia* (mugwort). Colour ranges have been set to 5 equal sized quantiles (i.e. each contains 20% of data). Note that absolute values for each taxon are different, see ranges on each colourbar. Images Crown Copyright, 2016, The Met Office. Based on digital spatial data licensed from the Centre for Ecology & Hydrology, copyright NERC (CEH).




Figure 4: Maps showing location of tree taxa in England and Wales. Units are number of trees per 1 km × 1 km grid square. Colour ranges have been set to 5 equal sized quantiles (i.e. each contains 20% of data). Note that absolute values for each taxon are different, see ranges on each colourbar. Tree locations of *Top left Alnus* (alder); *Top right Salix* (willow); *Bottom left Quercus* (oak); *Bottom right Fraxinus* (ash). Images Crown Copyright, 2016, The Met Office. Based on data licensed from National Tree Map, Copyright Bluesky International Limited, 2016; digital spatial data licensed from the Centre for Ecology & Hydrology, copyright NERC (CEH); and contains, or is based on, information supplied by the Forestry Commission.




Figure 5: Maps showing location of tree taxa in England and Wales. Units are number of trees per 1 km \times 1 km grid square. Colour ranges have been set to 5 equal sized quantiles (i.e. each contains 20% of data). Note that absolute values for each taxon are different, see ranges on each colourbar. Tree locations of *Top left Betula* (birch); *Top right Corylus* (hazel); *Bottom Pinus* (pine). Images Crown Copyright, 2016, The Met Office. Based on data licensed from National Tree Map, Copyright Bluesky International Limited, 2016; digital spatial data licensed from the Centre for Ecology & Hydrology, copyright NERC (CEH); and contains, or is based on, information supplied by the Forestry Commission.

336 The weed location maps in London show that dock is not nearly as abundant in the central area, while the other
337 weeds are more highly concentrated in the city. The grass map shows that there is less grass in central London, with
338 mixed values in the surrounding area. The tree maps show more variation between taxa. Notably pine stands out as
339 there is very little pine found in London, and while much higher values are seen in the surrounding area. Willow and
340 alder, are similar again, with fairly even levels of number of these taxa across the area. In the pine, hazel, birch and
341 oak maps, a peak can be seen south of the river just west of the centre of the map. This is a large green space in the
342 city (Richmond Park), containing many trees, and it shows up in these maps as an area with higher density of these
343 taxa.

344 4. Discussion

345 These maps of key allergenic tree, grass and weed taxa detail where these plants are located in the UK. They show
346 the different geographical distributions of the 12 taxa of trees, weeds and grass, and can be used to study plants in the
347 UK that are associated with allergy and allergic asthma. These have been made to study environmental exposure and
348 human health, although these new vegetation maps have many possible applications. Because the taxa mapped here
349 are all monitored by the UK pollen network, this would also allow further studies to compare to observational pollen
350 count data.

351 This novel method not only provides maps of many different plant types, but also at high resolution across England
352 and Wales, and Great Britain, and uniquely presents 12 key plant taxa using a consistent methodology.

353 We note that certain taxa are considered more important aeroallergens than others mapped in this study. We
354 highlight that the review by de Weger et al. (2012) found the following of our selected taxa highly allergenic: *Corylus*
355 (hazel), *Alnus* (alder), *Betula* (birch), *Platanus* (plane) and *Quercus* (oak), *Poaceae* (bulk grass, not species specific),
356 *Artemisia* (mugwort). Of the remaining taxa that we have mapped, we note that *Pinus* (pine) is not considered to
357 be allergenic. Neither is *Urtica* (nettle), although it has pollen which cannot be distinguished microscopically from
358 *Parietaria judaica* (de Weger et al., 2012) which is a potential aeroallergen and is frequently recorded in southern
359 Britain. Thus care would need to be taken if comparing these maps to observational pollen counts that *Parietaria*
360 *judaica* pollen grains were not being misclassified as *Urtica* (nettle). *Plantago* (plantain) is thought to be only a
361 minor cause of hay fever in Europe and has some cross-reactivity with grass pollen (de Weger et al., 2012). Similarly,
362 *Rumex* (dock, sorrels) is not considered to be important aeroallergen. *Salix* (willow) is both wind pollinated and insect
363 pollinated, it has minimal importance as an aeroallergen, with the airborne pollen loads often low.

364 We also note that [some sites in](#) the pollen network does also keep a watch out for other pollen taxa / species
365 including *Ambrosia* (ragweed), due to it being very highly allergenic (Dahl et al., 1999). However, it is not a native
366 plant nor currently commonly found in the UK, although it is highly invasive and there is a risk it may spread to the
367 UK in future with changes to the climate and land use (Ziska et al., 2011; Smith et al., 2013; Hamaoui-Laguel et al.,
368 2015; Pashley et al., 2015). As with other pollen, ragweed pollen has been shown to be transported long distances in




Figure 6: Maps showing location of weed taxa and grass in greater London, UK. *Top left*: percentage cover of *Rumex* (dock, sorrels); *Top right*: percentage cover of *Artemisia* (mugwort); *Centre left*: percentage cover of *Plantago* (plantain); *Centre right*: percentage cover of *Urtica* (nettle); *Bottom*: percentage cover of *Poaceae* (grass). Images Crown Copyright, 2016, The Met Office. Based on digital spatial data licensed from the Centre for Ecology & Hydrology, copyright NERC (CEH); and contains OS data Crown copyright (2016).




Figure 7: Maps showing location of tree taxa in greater London, UK. Units are number of trees per 1 km \times 1 km grid square. *Top left:* percentage cover of *Alnus* (alder); *Top right:* percentage cover of *Fraxinus* (ash); *Second row, left:* percentage cover of *Betula* (birch); *right:* percentage cover of *Corylus* (hazel); *Third row, left:* percentage cover of *Quercus* (oak); *right:* percentage cover of *Pinus* (pine); *Bottom:* percentage cover of *Salix* (willow). Images Crown Copyright, 2016, The Met Office. Based on data licensed from National Tree Map, Copyright Bluesky International Limited, 2016; digital spatial data licensed from the Centre for Ecology & Hydrology, copyright NERC (CEH); contains, or is based on, information supplied by the Forestry Commission; and contains OS data Crown copyright (2016).

369 the wind, and so areas with high concentrations of ragweed pollen in the air are more extensive than those where most
370 ragweed plants are located (Prank et al., 2013; Grewling et al., 2016; de Weger et al., 2016).

371 These maps fill a need for spatially detailed vegetation mapping of allergenic plants across a country, the first
372 maps of their kind in the UK. Moreover, the method presented here could be duplicated for other locations around the
373 world, with an initial assessment of data availability and expertise on plant types and land cover.

374 4.1. Uncertainty and comparisons with other results

375 Validation of inventories is generally not possible as they must include all data before they can be considered
376 inventories. Common practice is therefore to discuss the main elements of uncertainty and compare the inventory
377 results with relevant data including previous inventories (e.g. Skjøth et al. (2008) **Simpson1999, Oderbolz et al,
378 2013**). We have followed the same procedure by including these aspects in the following two main subsections:

379 4.1.1. Uncertainty with respect to the input data

380 The Land Cover Map 2007 dataset, and the National Forestry Inventory (NFI) dataset are both widely used envi-
381 ronmental datasets, appearing in many other peer reviewed studies. Nevertheless they contain some limitations that
382 will affect some of the species. The LCM2007 data set is a remote sensing product with 20-30 m resolutions used as
383 input from several satellites such as Landsat and SPOT. A consequence is smaller units will not be mapped. This is
384 however of minor importance for most of the weeds as we have applied an ecosystem approach combined with expert
385 estimates to estimate the abundance of all weeds. However it will have an some impact on *Artemisia* as this species is
386 often found along roads and rivers (e.g. **Essl et al, 2009**). This weed will therefore have a larger uncertainty than
387 the others. LCM will also have an impact on the mapping of grass areas in urban areas as grass ecosystems often are
388 associated with road sides (Skjøth et al., 2013), that cannot be mapped using the LCM2007. Furthermore, it is known
389 that grasses are also found in croplands under rotation (e.g. timothy and rye grass) that were not included in this
390 study. These areas are found most abundantly in the eastern parts of England, where the mapping suggest to contain
391 low amounts of grass areas. The Trees in Towns II dataset was used for urban taxa fractions and is the result of a
392 very thorough study of urban trees across many towns and cities. However the information was only available as one
393 nation-wide statistical data set and public data for London suggests, that not all tree species are uniformly distributed
394 within the towns. However a UK study comparing the UK towns Torbay, Wrexham, Glasgow and Edinburgh (**Rum-
395 ble et al, 2015**) suggests that tree species distribution within the towns are relatively similar, most likely caused by
396 a similar planting practice and land use type and, to a smaller degree, climate. This indicates that urban data in this
397 study is relatively robust, while in-town variations as seen over London are more uncertain. Finally, the Bluesky NTM
398 datasets high resolution (individual tree level resolution) and mapping methodology means that we have confidence
399 in the gridded tree count we created from it. Bluesky quote that “Independent verification shows that NTM identifies
400 at least 95% of tree canopy”. This suggests that the loss of tree information is below 5% nationally as well as within
401 each 1km grid cell. This accuracy is much higher than other more coarse data satellite based landcover products (e.g.

402 LCM2007, Corine Land Cover (CLC2000) or Globcover), that have shown to vary with with up to 50% in tree cover
403 and sometimes contain a direct misplacement of the tree resources (Skjøth et al., 2015b), CLC2000, which is much
404 lower. Hazel is in this respect a special case. Hazel is often found as as a bush or shrub in urban gardens and as an
405 important part in hedgerows. Neither of these sources can be expected to be mapped by the Bluesky data set due to
406 the 3 m limitation. Despite this limitation, then the application of the Bluesky data set has added a unique accuracy
407 to the tree resource, the urban areas in particular as the urban tree resource has never been mapped before within an
408 entire country.

409 4.1.2. *Uncertainty with respect to species information and inventory method*

410 The regional-level species information is what is likely to cause the largest uncertainties in our maps, here divided
411 into 13 different regions. These large geographical regions are not uniform in their distribution of species, but the
412 information is provided at this coarse scale, and as such this limits the quality of the species-level information within
413 these regions. Although the high-resolution tree database does mean that the locations of trees within these regions
414 is highly accurate, the exact distribution of the species on a fine scale is less certain. Previous inventories of tree
415 species information over the UK included up to 65 regions (Skjøth et al., 2008) showing that distribution of important
416 species like birch is highly uneven in the UK with hotspots found in a few regions in South England (Surrey), central
417 England (west Midlands) and North England (e.g Cumbria), respectively. A consequence is that some regions in our
418 inventory will then have a lower quantity of birches than reality, while other regions will have a large quantity than
419 reality. However comparing with the maps in Skjøth et al. (2008) this seems to be mainly an issue for birch causing
420 additional uncertainties with more than a factor of two, while oak, and alders have a much more uniform distribution,
421 thereby not being affected by this uncertainty. Outside the main forests oak and ash trees are known to completely
422 dominate the small woodlands. A consequence as that our mapping will underestimate the abundance of these two
423 species in forest sparse regions such as the eastern parts of England, while the more forest dense regions will not be
424 affected. With respect to total numbers within all of UK it has previously been estimated that this error is only going
425 to have a minor importance for oak (Skjøth et al., 2015b), while the impact on ash remains to be investigated. The
426 use of national specialist in assessing weed abundance has with success been used for ragweed throughout Europe
427 (Thibaudon et al., 2014; Karrer et al., 2015). Here this approach has been expanded to cover several specialists, each
428 with their independent assessment. The main uncertainty in this element is likely to be the number of specialists,
429 where a larger number of specialists could provide a more detailed picture. This is also related to the input data, as
430 abundance of some species are related to land use (e.g. nettle and *Artemisia*) such as fertilization. The LCM2007 does
431 not include land use but only land cover. Including land use could therefore improve the information for weeds and at
432 the same time potentially also improve the information on grasses.

433 The maps over London for alder, ash, birch, hazel, oak, pine and willow (Figure 7) appear to be almost 100%
434 spatially correlated. This is related to the fact that the two main data sets for the urban mapping are the urban tree
435 statistics (where there is only one set of nation-wide numbers) and GPS coordinates for trees that do not distinguish

436 between broadleaves/conifers or individual tree species. A comparison with the public road trees data set of
437 London that contain 716406 tree locations provides following information (alder=10325, ash=40870, birch=34839,
438 hazel=5597, oak=36853, pine=6399, willow=9102, other=572421), which provides a different overall distribution
439 than Figure 7 (e.g. oak being more abundant than birch). This suggests that the urban maps for individual cities or
440 towns should be treated with caution while the overall national distribution according to **Rumble et al (2015)**
441 must be assumed to be alright. Improvements in this area could be obtained with nation-wide high resolution satellite
442 products that can distinguish between broadleaved and conifers and with town specific tree species information. Some
443 of the allergenic species found in this inventory (e.g. hazel and mugwort) have never been mapped before. Comparison
444 with previous studies is therefore not possible. However a comparison of the mapping of oak, alder, birch and ash
445 with studies by Skjøth et al. (2015b) and Maskell et al. (2013) suggest that the patterns found throughout UK in this
446 study is similar to previous studies. This suggest that hazel, despite the limitations on not mapping hedgerows and
447 shrubs, will have similar quality as the other trees, thereby being unique not only for UK but the first of its kind in
448 Europe.

449 4.2. *Relevance to public health*

450 The maps presented here do not provide a forecast of pollen in the air at any one time, or human exposure, but
451 do provide the most likely locations of grass, tree and weed taxa that release allergenic pollen in Great Britain. The
452 allergenic taxa maps could provide information to local authorities and healthcare practitioners. They could be a useful
453 risk assessment tool to inform treatment or self-management of patients with long-term health conditions caused by
454 pollen allergy.

455 Hay fever and pollen-related asthma are only invoked by airborne pollen from allergenic plant taxa. To consider
456 the impact on human health due to exposure of the pollen grains, one must consider the timing of the pollen release,
457 and its dispersal. Along with the potential impact of climate change, air quality, changing allergenicity of pollen,
458 and possible mitigation options and their impact on public health, we discuss these important considerations in the
459 following sections.

460 4.2.1. *Timing of release of pollen grains*

461 A summary of the timing of peak pollen emissions (month of peak emissions) and emission strength categorised
462 by peak pollen count (number of grains per day) are given in Table 5. Results were calculated as averages of the day
463 of the year of peak emissions (presented by month) and peak pollen count (presented as categories) for each of the
464 species/taxa and across 9 sites in the UK pollen sampling network for the period 2006-2015. The categories for peak
465 pollen count (in units of number of grains per day) are as follows: Low = 0-10; Low-Med = 10-50; Med = 50-100;
466 Med-High = 100-300; High = 300-500. The sites span a wide spatial distribution (from south to north - Plymouth, Isle
467 of Wight, London Islington, Cardiff, Worcester, Cambridge, Leicester, Belfast and Invergowrie). As there were many
468 missing data, the years and sites used for these calculations are listed for each species/taxa. Note that for *Corylus*

taxa or species	common name	years and sites with data available	month of peak emissions	emissions strength*
<i>Corylus avellana</i>	common hazel	2006-12 ^W	February	Med
<i>Alnus</i>	alder	2006-12 ^W	February/March	Med-High
<i>Betula</i>	birch	2006-15 ^{P,IoW,Cf,Ig} 2006-12 ^W 2006-11 ^{LI} 2006, 2008, 2011-15 ^L 2006-8, 2012 ^B 2006-14 ^{Cb}	April	High
<i>Fraxinus</i>	ash	2006-15 ^{P,IoW,Cf,Ig} 2006-12 ^W 2006-11 ^{LI} 2006, 2008, 2011-15 ^L 2006-9, 2012 ^B 2006-14 ^{Cb}	April	Med-High
<i>Salix</i>	willow	2006-15 ^{P,IoW,Cf,Ig} 2006-12 ^W 2006-11 ^{LI} 2008, 2011-15 ^L 2006-9, 2012 ^B 2006-14 ^{Cb}	April	Low-Med
<i>Pinus</i>	pine	2008 ^L	May	Med-High
<i>Quercus</i>	oak	2006-15 ^{P,IoW,Cf,Ig} 2006-12 ^W 2006-11 ^{LI} 2006, 2008, 2011-15 ^L 2006-9, 2011-12 ^B 2006-14 ^{Cb}	May	Med-High
<i>Poaceae</i>	grass	2006-15 ^{P,IoW,Cf,Ig,W} 2006-11 ^{LI} 2006, 2008-15 ^L 2006-9, 2011-12 ^B 2006-14 ^{Cb}	June	Med-High
<i>Rumex</i>	dock, sorrel	2008 ^L	June	Low-Med
<i>Urtica</i>	nettle	2006-15 ^{P,IoW,Cf,Ig} 2006-12 ^W 2006-11 ^{LI} 2006, 2008, 2011-15 ^L 2006-9, 2011-12 ^B 2006-14 ^{Cb}	June/July	High
<i>Plantago</i>	plantain	2008 ^L	June	Low
<i>Artemisia</i>	mugwort	2006-15 ^{P,IoW} 2009, 2011-15 ^{Cf} 2007, 2009, 2015 ^{Ig} 2006-12 ^W 2006-11 ^{LI} 2006, 2008, 2011-15 ^L 2006-7, 2009, 2011 ^B 2006-14 ^{Cb}	July	Low

Table 5: A summary of the timing of peak pollen emissions for each of the species/taxa and across 9 sites in the UK pollen sampling network for the period 2006-2015. Month of peak emissions gives the timing of peak pollen emissions; while emission strength is categorised* by peak pollen count (number of grains per day). As there were many missing data, the years and sites used for these calculations are listed for each species/taxa. The sites span a wide spatial distribution (from south to north - Plymouth^P, Isle of Wight^{IoW}, London Islington^{LI}, Cardiff^{Cf}, Worcester^W, Cambridge^{Cb}, Leicester^L, Belfast^B, Invergowrie^{Ig}). *Based on peak daily pollen count (in units of number of grains per day): Low = 0-10, Low-Med = 10-50, Med = 50-100, Med-High = 100-300, High = 300-500.

469 *avellana* and *Alnus* only the Worcester data were used as this was the only site with data available before March,
470 when pollen emissions for these species are likely in many locations. Note that sensitivity tests on a range of different
471 time periods have been made and the same general trends were evident.

472 Pollen grains contain the male reproductive cells of seed plants, and the timing of their development and release
473 varies depending on the taxon and environmental conditions, particularly meteorology (Emberlin et al., 2007). The
474 production of pollen is dependent not only on the current meteorological conditions (including day length, temper-
475 ature, precipitation, and wind speed/direction), but also on the conditions and water availability experienced in the
476 year prior during which pollen is formed (Emberlin et al., 2007). Land management is also critical, for example if
477 fields are cut or grazed by animals then the potential to release pollen is significantly affected. Any changes in these
478 conditions affect the phenology of the tree and thus the timing of the onset of pollen release, the total volume of pollen
479 produced, and the length of the flowering season (Dahl et al., 2012). Typically in the UK, the pollen season for tree
480 taxa ranges from January to April for early flowering taxa such as alder and hazel, to mid-March to end of July for the
481 later taxa, including oak and pine. Grass flowering occurs generally between May and early-September, coincident
482 with many taxa of weeds with allergenic pollen, see Worcester University (2017). Depending on the taxon, pollen
483 grains are dispersed by wind, animals or water.

484 4.2.2. *Dispersion of pollen grains*

485 Dispersion of pollen grains once they are released from a plant is dependent on many factors, predominantly
486 meteorological. Land management is also important, for example hay cutting can shed substantial amounts of pollen
487 in an acute manner. Pollen can be transported large distances from its source (Sofiev et al., 2012), with the main
488 factors in their dispersion being wind speed and direction, as well as precipitation (which brings down the pollen to
489 the ground). The distance that a pollen grain travels depends on the taxon of plant, as each of the pollen grains are
490 different sizes and shapes and so have different aerodynamic properties; and also on the release height of the grain
491 (for example tree pollen are released higher from the ground and are more likely to be transported long-range). Pollen
492 may travel some distance, and the concentration of pollen (particles per m³) is not just a localised phenomenon; it
493 varies on a fairly broad geographical scale. Pollen grain counts have been found to correlate across distances of 20 km
494 (Erbas et al., 2007) and 41 km (Pashley et al., 2009), with the potential for pollen to travel much further (Skjøth et al.,
495 2007), indeed across the globe (Sofiev et al., 2012).

496 4.2.3. *Effect of climate change*

497 The local effect of climate change on pollen production, release timing, transport and deposition is highly complex,
498 and its impact on pollen allergies is very uncertain (Osborne and Eggen, 2014). There is some evidence that climate
499 change may result in earlier seasonal appearance of respiratory symptoms and longer duration of exposure to pollen
500 (Fitter and Fitter, 2002; Vardoulakis and Heaviside, 2012). Across the Northern Hemisphere, early onset of spring
501 has been measured, with some tree taxa releasing pollen earlier (Emberlin et al., 1997; Frei, 1998; Emberlin et al.,

502 2002; Beggs, 2004) and cases of lengthened pollen season (Ziska et al., 2011). Further warming of global temperature
503 and increases in atmospheric CO₂ concentration may lead to great pollen release through increased plant productivity,
504 unless the plants are limited by other factors such as water stress. Failure to frost means that the pollen season may be
505 prolonged (Gezon et al., 2016). Climate change may also affect the allergenicity of pollen for some taxa (Cecchi et al.,
506 2010). Furthermore, the effect of climate change on the frequency or severity of thunderstorms, and thus the health
507 impact of thunderstorm asthma (Elliot et al., 2013) (where it is thought pollen plays a role) under a future climate is
508 very uncertain.

509 4.2.4. *Interaction with air quality*

510 People who live in urban areas have been shown to be more affected by pollen allergies (asthma and allergic
511 rhinitis) than those who live in rural areas (Ehrenstein et al., 2000; Riedler et al., 2001; D'Amato et al., 2007). Ur-
512 ban environments, with high levels of vehicle emissions have been shown to coincide with increased pollen-induced
513 respiratory allergies. While it has been known for some time that exposure to air pollution prior to pollen exposure
514 can exacerbate symptoms and lower the threshold of pollen required to trigger symptoms in allergy sufferers (Molfino
515 et al., 1991; Emberlin, 1998), further research is required to characterise the interaction between chemical air pol-
516 lutants and pollen. For example, to fully understand and quantify the effect of exposure to both allergenic pollen
517 and pollutants, the effect on both the allergenicity (such as increased allergenicity of pollen which had been exposed
518 to NO₂ found by (Cuinica et al., 2013)) and volume of pollen grains released under increased air pollution must be
519 determined. The health impacts of all these factors, in high co-exposure areas need to be considered. Co-exposure of
520 pollen and air pollutants (namely NO₂, SO₂ and particulate matter PM_{2.5} and PM₁₀ and ozone) is currently an active
521 area of research (Mücke et al. (2014); Ørby et al. (2015)); allergens from birch pollen have been found bound to
522 particulate matter, PM₁₀ which means the pollen allergen can travel into the low respiratory tract (Suring et al., 2016).
523 Vardoulakis et al. (2015) discuss how a range of health risks from indoor pollutants, including pollen, might change
524 under a changing climate. When considering the public health relevance of these taxa maps, it is important to note the
525 interaction between pollen and air quality, as well as the spatial variability of air pollution (e.g. see Defra / Met Office
526 air quality maps (Met Office, 2017a; Defra, 2017)).

527 4.2.5. *Changing allergenicity*

528 The allergen content of pollen has also been shown to vary throughout the season and across regions. Buters
529 et al. (2015) found that grass pollen allergen release (potency) varied significantly, and that differences were found per
530 location in Europe, and time in the pollen season. They concluded that although pollen count is a proxy for exposure,
531 it is not a measure of the actual allergens in the air.

532 4.2.6. *Mitigation and adaptation*

533 To mitigate the health effects of pollen from allergenic trees in urban areas, adaptation measures have been sug-
534 gested. Policies to oversee planting of allergenic taxa in urban areas, e.g. *Platanus* (London plane), and the use of

535 an allergenicity scale in the US (Seitz and Francisco Escobedo, 2009) to select suitable taxa for urban planting have
536 been proposed. Cecchi et al. (2010); D'Amato (2011) recommend planting non-allergenic trees such as *Palmaceae*
537 (palm) and *Ulmaceae* (elm), avoiding *Cupressaceae* (cypress), *Betulaceae* (birch) and *Oleaceae* (olive). *Cryptomeria*
538 *japonica* (Sugi, or Japanese cedars) were planted extensively in Japan, Asia and North America after the second world
539 war. These highly allergenic trees have had a large health effect on populations (Okuda, 2003; Yamada et al., 2014).
540 Furthermore, the current practice of the planting of solely male trees in urban areas in order to reduce street litter from
541 seeds and fruit increases the total pollen load, and so consideration of reduction of pollen exposure through female
542 tree planting could potentially mitigate effects. The services and disservices that a particular tree taxon and canopy
543 design may provide must be considered by local decision makers (Salmond et al., 2016).

544 Other vegetation management schemes such as grass cutting regimes where grass is cut before it flowers could be
545 used in areas where these maps show that there is a high coverage of grass close to high population centres. It should
546 be noted, however, that if plants are routinely managed to maintain vegetative phases of growth and restrict flowering
547 there will be consequent negative impacts on biodiversity.

548 **5. Conclusions**

549 In this paper we present a methodology to produce location maps of selected trees, grass and weeds, to study
550 plant taxa in the UK that are associated with allergy and allergic asthma. The taxa and species used in this study are
551 all monitored as part of the UK pollen monitoring network. These have been made in order to study environmental
552 exposure and human health, although these new vegetation maps have many possible applications.

553 The novel method used in this study uniquely maps 12 key plant taxa using a consistent, reproducible methodology,
554 covering more plant types, and higher spatial resolution across England and Wales, and Great Britain, than currently
555 available elsewhere.

556 To create these maps, several different observational and modelled datasets were combined. These include land
557 use datasets and tree taxa databases including those from the Forestry Commission and the National Tree Map from
558 Bluesky International. A framework was designed to combine these datasets allowing integration of existing UK land
559 use and vegetation datasets with the taxa data. We presented 1 km resolution maps of 12 taxa of trees, grass and weeds
560 found in the UK, and plots showing local level detail around the city of London.

561 The next step of this work is to combine these taxa maps with health data including emergency hospital admissions
562 for asthma to study the impact of different taxa in the UK. The maps could also be combined with population data in a
563 risk assessment framework. This will provide valuable information on which taxa in the UK have the most significant
564 health impact for asthma. Furthermore, in the future these maps could also form part of a UK pollen forecast model,
565 to provide detailed taxon-specific, localised information to the public.

566 Acknowledgements

567 Funding provided by the National Institute for Health Research Health Protection Research Unit (NIHR HPRU) in
568 Environmental Change and Health at the London School of Hygiene and Tropical Medicine in partnership with Public
569 Health England, and in collaboration with the University of Exeter, University College London, and the Met Office.
570 Additional financial support was provided by the Joint DECC/Defra Met Office Hadley Centre Climate Programme
571 (GA01101). Observed pollen data for Table 5 were collected by the following sites and we thank all the individual
572 collectors at each site for this: Belfast, Cambridge, Cardiff, Invergowrie, Isle of Wight, Leicester, London Islington,
573 Plymouth and Worcester. We also thank the Midlands Asthma and Allergy Research Association (MAARA) as they
574 support the collection of aerobiological data at Leicester; and the Met Office who have operated the UK pollen network
575 since 2011.

576 The authors thank: Phil Bentley for contribution to scripting in python for tree and grass mapping analysis, and
577 big data handling; Neil Kaye for contribution to tree mapping work and expertise in ArcGIS; Nick Morgan and Nick
578 Dobson for help with ArcGIS and data management; Yolanda Clewlow head of health research and services at the
579 Met Office, for managing the UK pollen network and establishing relationships with external health colleagues; Ben
580 Ditchburn (Forestry Commission) for initial discussions on tree mapping and forestry datasets. Furthermore, the
581 authors warmly thank the anonymous reviewer for their constructive comments which have improved the paper.

582 The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, the Department of
583 Health or Public Health England.

584 Appendix A. Supplementary material

585 Appendix A.1. Supplementary Figures

586 Adams-Groom, B., Emberlin, J., Corden, J., Millington, W., Mullins, J., 2002. Predicting the start of the birch pollen season at London, Derby and
587 Cardiff, United Kingdom, using a multiple regression model, based on data from 1987 to 1997. *Aerobiologia* 18 (2), 117–123.

588 URL <http://dx.doi.org/10.1023/A:1020698023134>

589 Asthma UK, 2017. Asthma UK, last accessed 06/03/2017.

590 URL <https://www.asthma.org.uk/advice/triggers/pollen/>

591 Bauchau, V., Durham, S., 2004. Prevalence and rate of diagnosis of allergic rhinitis in Europe. *European Respiratory Journal* 24 (5), 758–764.

592 URL <http://erj.ersjournals.com/content/24/5/758>

593 Beggs, P. J., 2004. Impacts of climate change on aeroallergens: past and future. *Clinical & Experimental Allergy* 34 (10), 1507–1513.

594 URL <http://dx.doi.org/10.1111/j.1365-2222.2004.02061.x>

595 Bluesky, National Tree Map, 2015. Bluesky National Tree Map. [Http://www.bluesky-world.com/#!national-tree-map/c1pqz](http://www.bluesky-world.com/#!national-tree-map/c1pqz), Accessed: 25/01/2016.

596 Bousquet, J., Van Cauwenberge, P., Khaltaev, N., Aria Workshop Group, World Health Organisation, 2001. Allergic rhinitis and its impact on
597 asthma. *The Journal of Allergy and Clinical Immunology* 108(Suppl.5), 147–334.

598 Bousquet, P.-J., Chinn, S., Janson, C., Kogevinas, M., Burney, P., Jarvis, D., 2007. Geographical variation in the prevalence of positive skin tests to
599 environmental aeroallergens in the European Community Respiratory Health Survey I. *Allergy* 62 (3), 301–309.

600 URL <http://dx.doi.org/10.1111/j.1398-9995.2006.01293.x>




Figure A.8: Maps showing location of four weeds in Great Britain, in units of percentage cover per $1 \text{ km} \times 1 \text{ km}$ grid box. *Top left*: percentage cover of *Rumex* (dock, sorrels); *Top right*: percentage cover of *Urtica* (nettle); *Bottom left*: percentage cover of *Plantago* (plantain); *Bottom right*: percentage cover of *Artemisia* (mugwort). Images Crown Copyright, 2016, The Met Office. Based on digital spatial data licensed from the Centre for Ecology & Hydrology, copyright NERC (CEH).




Figure A.9: Maps showing location of tree taxa in England and Wales. Units are number of trees per 1 km \times 1 km grid square. Note that absolute values for each taxon are different, see ranges on each colourbar. Tree locations of *Top left Alnus* (alder); *Top right Salix* (willow); *Bottom left Quercus* (oak); *Bottom right Fraxinus* (ash). Images Crown Copyright, 2016, The Met Office. Based on data licensed from National Tree Map, Copyright Bluesky International Limited, 2016; digital spatial data licensed from the Centre for Ecology & Hydrology, copyright NERC (CEH); and contains, or is based on, information supplied by the Forestry Commission.




Figure A.10: Maps showing location of tree taxa in England and Wales. Units are number of trees per 1 km \times 1 km grid square. Note that absolute values for each taxon are different, see ranges on each colourbar. Tree locations of *Top left Betula* (birch); *Top right Corylus* (hazel); *Bottom Pinus* (pine). Images Crown Copyright, 2016, The Met Office. Based on data licensed from National Tree Map, Copyright Bluesky International Limited, 2016; digital spatial data licensed from the Centre for Ecology & Hydrology, copyright NERC (CEH); and contains, or is based on, information supplied by the Forestry Commission.

601 Brewer, A., 2013. NFI Preliminary Report NFI preliminary estimates of quantities of broadleaved species in British woodlands, with special focus
602 on ash. Tech. rep., Forestry Commission, Great Britain.

603 Britt, C., Johnston, M., 2008. Trees in Towns II, A new survey of urban trees in England and their condition and management. Research for amenity
604 trees. Department for Communities and Local Government.

605 Brus, D. J., Hengeveld, G. M., Walvoort, D. J. J., Goedhart, P. W., Heidema, a. H., Nabuurs, G. J., Gunia, K., Apr. 2011. Statistical mapping of tree
606 species over Europe. *European Journal of Forest Research* 131 (1), 145–157.
607 URL <http://link.springer.com/10.1007/s10342-011-0513-5>

608 Burbach, G. J., Heinzerling, L. M., Edenharter, G., Bachert, C., Bindslev-Jensen, C., Bonini, S., Bousquet, J., Bousquet-Rouanet, L., Bousquet,
609 P. J., Bresciani, M., Bruno, A., Canonica, G. W., Darsow, U., Demoly, P., Durham, S., Fokkens, W. J., Giavi, S., Gjomarkaj, M., Gramiccioni, C.,
610 Haahtela, T., Kowalski, M. L., Magyar, P., Muraközi, G., Orosz, M., Papadopoulos, N. G., Röhnelt, C., Stingl, G., Todo-Bom, A., Von Mutius,
611 E., Wiesner, A., Wöhrl, S., Zuberbier, T., 2009. GA2LEN skin test study II: clinical relevance of inhalant allergen sensitizations in Europe.
612 *Allergy* 64 (10), 1507–1515.
613 URL <http://dx.doi.org/10.1111/j.1398-9995.2009.02089.x>

614 Buters, J., Prank, M., Sofiev, M., Pusch, G., Albertini, R., Annesi-Maesano, I., Antunes, C., Behrendt, H., Berger, U., Brandao, R., Celenk, S.,
615 Galan, C., Grewling, L., Jackowiak, B., Kennedy, R., Rantio-Lehtimäki, A., Reese, G., Sauliene, I., Smith, M., Thibaudon, M., Weber, B.,
616 Cecchi, L., 2015. Variation of the group 5 grass pollen allergen content of airborne pollen in relation to geographic location and time in season.
617 *Journal of Allergy and Clinical Immunology* 136 (1), 87 – 95.e6.
618 URL <http://www.sciencedirect.com/science/article/pii/S0091674915004121>

619 Cecchi, L., 2012. Introduction. In: Sofiev, M., Bergmann, K.-C. (Eds.), *Allergenic pollen: a review of the production, release, distribution and*
620 *health impacts*. Springer, Ch. 1, pp. 1–7.

621 Cecchi, L., D’Amato, G., Ayres, J. G., Galan, C., Forastiere, F., Forsberg, B., Gerritsen, J., Nunes, C., Behrendt, H., Akdis, C., Dahl, R., Annesi-
622 Maesano, I., 2010. Projections of the effects of climate change on allergic asthma: the contribution of aerobiology. *Allergy* 65 (9), 1073–1081.

623 Centre for Ecology & Hydrology, UK, July 2011. Land Cover Map 2007 dataset documentation, countryside survey technical report. Tech. rep.,
624 NERC/Centre for Ecology & Hydrology.

625 Corine, 2017. Corine, last accessed 06/03/2017.
626 URL <http://land.copernicus.eu/pan-european/corine-land-cover>

627 Csornai, G., Mikus, G., Nádor, G., Hubik, I., László, I., Suba, Z., 2011. The first seven years of the remote sensing based ragweed monitoring and
628 control system. *EARSeL eProceedings* 10(2), 110–118.

629 Cuinica, L. G., Abreu, I., Esteves da Silva, J., Dec. 2013. Effect of air pollutant NO₂ on *Betula pendula*, *Ostrya carpinifolia* and *Carpinus betulus*
630 pollen fertility and human allergenicity. *Environmental pollution (Barking, Essex : 1987)* 186C (2), 50–55.
631 URL <http://www.ncbi.nlm.nih.gov/pubmed/24361564>

632 Dahl, Å., Galan, C., Hajkova, L., Pauling, A., Sikoparija, B., Smith, M., Vokou, D., 2012. The onset, course and intensity of the pollen season. In:
633 Sofiev, M., Bergmann, K.-C. (Eds.), *Allergenic pollen: a review of the production, release, distribution and health impacts*. Springer, Ch. 3, pp.
634 29–70.

635 Dahl, Å., Strandhede, S.-O., Wihl, J.-Å., 1999. Ragweed – An allergy risk in Sweden? *Aerobiologia* 15 (4), 293–297.
636 URL <http://dx.doi.org/10.1023/A:1007678107552>

637 D’Amato, G., 2011. Effects of climatic changes and urban air pollution on the rising trends of respiratory allergy and asthma. *Multidisciplinary*
638 *Respiratory Medicine* 6 (1), 1–10.
639 URL <http://dx.doi.org/10.1186/2049-6958-6-1-28>

640 D’Amato, G., Cecchi, L., Bonini, S., Nunes, C., Annesi-Maesano, I., Behrendt, H., Liccardi, G., Popov, T., Van Cauwenberge, P., 2007. Allergenic
641 pollen and pollen allergy in Europe. *Allergy* 62 (9), 976–990.
642 URL <http://dx.doi.org/10.1111/j.1398-9995.2007.01393.x>

643 de Weger, L., Bergmann, K.-C., Rantio-Lehtimäki, A., Dahl, Å., Buters, J., Déchamp, C., Belmonte, J., Thibaudon, M., Cecchi, L., Besancenot,

644 J.-P., Galán, C., Waisel, Y., 2012. The onset, course and intensity of the pollen season. In: Sofiev, M., Bergmann, K.-C. (Eds.), Allergenic pollen:
645 a review of the production, release, distribution and health impacts. Springer, Ch. 6, pp. 161–216.

646 de Weger, L. A., Pashley, C. H., Šikoparija, B., Skjøth, C. A., Kasprzyk, I., Grewling, L., Thibaudon, M., Magyar, D., Smith, M., 2016. The
647 long distance transport of airborne Ambrosia pollen to the UK and the Netherlands from Central and south Europe. *International Journal of*
648 *Biometeorology* 60 (12), 1829–1839.
649 URL <http://dx.doi.org/10.1007/s00484-016-1170-7>

650 Defra, 2017. Defra daily air quality index, last accessed 06/03/2017.
651 URL <http://uk-air.defra.gov.uk/air-pollution/daq>

652 Ehrenstein, V., Mutius, V., Illi, Baumann, Bhm, Kries, V., 2000. Reduced risk of hay fever and asthma among children of farmers. *Clinical &*
653 *Experimental Allergy* 30 (2), 187–193.
654 URL <http://dx.doi.org/10.1046/j.1365-2222.2000.00801.x>

655 Elliot, A. J., Hughes, H. E., Hughes, T. C., Locker, T. E., Brown, R., Sarran, C., Clewlow, Y., Murray, V., Bone, A., Catchpole, M., McCloskey, B.,
656 Smith, G. E., Oct. 2013. The impact of thunderstorm asthma on emergency department attendances across London during July 2013. *Emergency*
657 *medicine journal : EMJ, emermed-2013-203122-*.
658 URL <http://emj.bmj.com/content/early/2013/10/08/emered-2013-203122.abstract>

659 Emberlin, J., 1998. The effects of air pollution on allergenic pollen. Vol. 8, 53. pp. 164—167.

660 Emberlin, J., Detandt, M., Gehrig, R., Jaeger, S., Nolard, N., Rantio-Lehtimäki, A., Sep. 2002. Responses in the start of Betula (birch) pollen
661 seasons to recent changes in spring temperatures across Europe. *International journal of biometeorology* 46 (4), 159–70.
662 URL <http://www.ncbi.nlm.nih.gov/pubmed/12242471>

663 Emberlin, J., Mullins, J., Corden, J., Millington, W., Brooke, M., Savage, M., Jones, S., Jan. 1997. The trend to earlier birch pollen seasons in the
664 U.K.: A biotic response to changes in weather conditions? *Grana* 36 (1), 29–33.
665 URL <http://dx.doi.org/10.1080/00173139709362586>

666 Emberlin, J., Smith, M., Close, R., Adams-Groom, B., Jan. 2007. Changes in the pollen seasons of the early flowering trees *Alnus* spp. and *Corylus*
667 spp. in Worcester, United Kingdom, 1996-2005. *International Journal of Biometeorology* 51 (3), 181–91.
668 URL <http://www.ncbi.nlm.nih.gov/pubmed/17024396>

669 Erbas, B., Chang, J.-H., Dharmage, S., Ong, E. K., Hyndman, R., Newbiggin, E., Abramson, M., 2007. Do levels of airborne grass pollen influence
670 asthma hospital admissions?. *Clinical & Experimental Allergy* 37 (11), 1641 – 1647.

671 ESA, 2017. ESA Landcover map, last accessed 06/03/2017.
672 URL <http://www.esa-landcover-cci.org/>

673 ESRI, 2017. ESRI, Redlands, US, last accessed 06/03/2017.
674 URL <http://www.esri.com/software/arcgis>

675 Fitter, A. H., Fitter, R. S. R., 2002. Rapid changes in flowering time in british plants. *Science* 296 (5573), 1689 – 1692.

676 Forestry Commission, Great Britain, 2011. NFI 2011 woodland map GB. Tech. rep.

677 Forestry Commission, Great Britain, 2012. Standing timber volume for coniferous trees in Britain: National Forest Inventory report. Tech. rep.,
678 Forestry Commission, Great Britain.

679 Frei, T., 1998. The effects of climate change in Switzerland 1969-1996 on airborne pollen quantities from hazel, birch and grass. *Grana* 37 (3),
680 172–179.
681 URL <http://dx.doi.org/10.1080/00173139809362662>

682 Gezon, Z. J., Inouye, D. W., Irwin, R. E., 2016. Phenological change in a spring ephemeral: implications for pollination and plant reproduction.
683 *Global Change Biology*, n/a–n/a.
684 URL <http://dx.doi.org/10.1111/gcb.13209>

685 Greiner, A. N., Hellings, P. W., Rotiroti, G., Scadding, G. K., 2012. Allergic rhinitis. *The Lancet* 378 (9809), 2112 – 2122.
686 URL <http://www.sciencedirect.com/science/article/pii/S014067361160130X>

687 Grell, G., Freitas, S. R., Stuefer, M., Fast, J., 2011. Inclusion of biomass burning in WRF-Chem: impact of wildfires on weather forecasts.
688 Atmospheric Chemistry and Physics 11 (11), 5289–5303.
689 URL <http://www.atmos-chem-phys.net/11/5289/2011/>

690 Grewling, L., Bogawski, P., Jenerowicz, D., Czarnecka-Operacz, M., Šikoparija, B., Skjøth, C. A., Smith, M., 2016. Mesoscale atmospheric
691 transport of ragweed pollen allergens from infected to uninfected areas. International Journal of Biometeorology, 1–8.
692 URL <http://dx.doi.org/10.1007/s00484-016-1139-6>

693 Grewling, L., Šikoparija, B., Skjøth, C., Radišić, P., Apatini, D., Magyar, D., Páldy, A., Yankova, R., Sommer, J., Kasprzyk, I., Myszkowska, D.,
694 Uruska, A., Zimny, M., Puc, M., Jäger, S., Smith, M., 2012. Variation in Artemisia pollen seasons in Central and Eastern Europe. Agricultural
695 and Forest Meteorology 160, 48 – 59.
696 URL <http://www.sciencedirect.com/science/article/pii/S0168192312000937>

697 Hamaoui-Laguél, L., Vautard, R., Liu, L., Solmon, F., Viovy, N., Khvorostyanov, D., Essl, F., Chuine, I., Colette, A., Semenov, M. A., Schaffhauser,
698 A., Storkey, J., Thibaudon, M., Epstein, M. M., 2015. Effects of climate change and seed dispersal on airborne ragweed pollen loads in europe.
699 Nature Climate Change 5 (8), 766 – 771.

700 IGBP, 2017. IGBP, last accessed 06/03/2017.
701 URL <http://glcf.umd.edu/data/lc/>

702 Karrer, G., Skjøth, C., Šikoparija, B., Smith, M., Berger, U., Essl, F., 2015. Ragweed (ambrosia) pollen source inventory for Austria. Science of
703 The Total Environment 523, 120 – 128.
704 URL <http://www.sciencedirect.com/science/article/pii/S0048969715003927>

705 Kaye, N., Burgess, P., 2013. A proposed methodology for quantifying and mapping habitat resilience in devon. Tech. rep., Met Office Hadley
706 Centre, <http://www.metoffice.gov.uk/hadobs/climviz/Culm/>.

707 Maskell, L., Henrys, P., Norton, L., Smart, S., Wood, C., January 2013. Distribution of Ash trees (*Fraxinus excelsior*) in Countryside Survey data.
708 Tech. rep.

709 Met Office, 2017a. Met Office air quality, last accessed 06/03/2017.
710 URL <http://www.metoffice.gov.uk/guide/weather/air-quality>

711 Met Office, 2017b. Met Office pollen forecast, last accessed 06/03/2017.
712 URL <http://www.metoffice.gov.uk/health/public/pollen-forecast>

713 MODIS Vegetation Continuous Fields, 2017. MODIS Vegetation Continuous Fields, last accessed 06/03/2017.
714 URL <http://glcf.umd.edu/data/vcf/>

715 Molfino, N., Wright, S., Katz, I., Tarlo, S., Silverman, F., McClean, P., Slutsky, A., Zamel, N., Szalai, J., Raizenne, M., 1991. Originally published
716 as volume 2, issue 8761 effect of low concentrations of ozone on inhaled allergen responses in asthmatic subjects. The Lancet 338 (8761), 199
717 – 203.
718 URL <http://www.sciencedirect.com/science/article/pii/014067369190346Q>

719 Morton, D., Rowland, C., Wood, C., Meek, L., Marston, C., Smith, G., Wadsworth, R., Simpson, I., July 2011. Final report for LCM2007 - the new
720 UK land cover map. countryside survey technical report no 11/07. Tech. rep., NERC/Centre for Ecology & Hydrology.
721 URL <http://nora.nerc.ac.uk/14854/>

722 Mücke, H.-G., Wagener, S., Werchan, M., Bergmann, K.-C., 2014. Measurements of particulate matter and pollen in the city of Berlin. Urban
723 Climate 10, Part 4, 621 – 629.
724 URL <http://www.sciencedirect.com/science/article/pii/S2212095514000236>

725 Netuveli, G., Hurwitz, B., Levy, M., Fletcher, M., Barnes, G., Durham, S. R., Sheikh, A., 2005. Ethnic variations in UK asthma frequency,
726 morbidity, and health-service use: a systematic review and meta-analysis. The Lancet 365 (9456), 312–317.
727 URL [http://dx.doi.org/10.1016/S0140-6736\(05\)17785-X](http://dx.doi.org/10.1016/S0140-6736(05)17785-X)

728 Newson, R. B., van Ree, R., Forsberg, B., Janson, C., Lötvall, J., Dahlén, S.-E., Toskala, E. M., Bælum, J., Brożek, G. M., Kasper, L., Kowalski,
729 M. L., Howarth, P. H., Fokkens, W. J., Bachert, C., Keil, T., Krämer, U., Bislimovska, J., Gjomarkaj, M., Loureiro, C., Burney, P. G. J., Jarvis,

730 D., 2014. Geographical variation in the prevalence of sensitization to common aeroallergens in adults: the GA2LEN survey. *Allergy* 69 (5),
731 643–651.
732 URL <http://dx.doi.org/10.1111/all.12397>

733 NHS Choices, 2017. NHS Choices, hay-fever, last accessed 06/03/2017.
734 URL <http://www.nhs.uk/conditions/Hay-fever/Pages/Introduction.aspx>

735 Office of National Statistics, 2017. Office of National Statistics, last accessed 06/03/2017.
736 URL <https://www.ons.gov.uk/>

737 O'Hagan, A., Buck, C. E., Daneshkhan, A., Eiser, J. R., Garthwaite, P. H., Jenkinson, D. J., Oakley, J. E., Rakow, T., 2006. *Uncertain Judgements:
738 Eliciting Experts' Probabilities*. John Wiley & Sons, Ltd.
739 URL <http://dx.doi.org/10.1002/0470033312.fmatter>

740 Okuda, M., 2003. Epidemiology of japanese cedar pollinosis throughout japan. *Annals of Allergy, Asthma & Immunology* 91 (3), 288 – 296.
741 URL <http://www.sciencedirect.com/science/article/pii/S1081120610635326>

742 Ørby, P., Peel, R., Skjøth, C., Schlünssen, V., Bønløkke, J., Ellermann, T., Brændholt, A., Sigsgaard, T., Hertel, O., Jan. 2015. An assessment of
743 the potential for co-exposure to allergenic pollen and air pollution in Copenhagen, Denmark. *Urban Climate*.
744 URL <http://www.sciencedirect.com/science/article/pii/S2212095514001102>

745 Osborne, N., Eggen, B., 2014. Pollen and asthma: impacts of anthropogenic climate change. *Living With Environmental Change*, LWEC, NERC.

746 Pashley, C., Fairs, A., Edwards, R., Bailey, J., Corden, J., Wardlaw, A., 2009. Reproducibility between counts of airborne allergenic pollen from
747 two cities in the east midlands, uk. *Aerobiologia* 25 (4), 249 – 263.

748 Pashley, C. H., Satchwell, J., Edwards, R. E., 2015. Ragweed pollen: is climate change creating a new aeroallergen problem in the UK? *Clinical &
749 Experimental Allergy* 45 (7), 1262–1265.
750 URL <http://dx.doi.org/10.1111/cea.12572>

751 Prank, M., Chapman, D. S., Bullock, J. M., Belmonte, J., Berger, U., Dahl, Å., Jager, S., Kovtunen, I., Magyar, D., Niemela, S., Rantio-
752 Lehtimäki, A., Rodinkova, V., Sauliene, I., Severova, E., Sikoparija, B., Sofiev, M., 2013. An operational model for forecasting ragweed pollen
753 release and dispersion in Europe. *Agricultural and Forest Meteorology* 182-183, 43 – 53.
754 URL <http://www.sciencedirect.com/science/article/pii/S0168192313002104>

755 Python Software Foundation, 2017. Python Software Foundation, last accessed 06/03/2017.
756 URL <https://www.python.org/>

757 Riedler, J., Braun-Fahrlander, C., Eder, W., Schreuer, M., Waser, M., Maisch, S., Carr, D., Schierl, R., Nowak, D., von Mutius, E., 2001. Exposure
758 to farming in early life and development of asthma and allergy: a cross-sectional survey. *The Lancet* 358 (9288), 1129 – 1133.
759 URL <http://www.sciencedirect.com/science/article/pii/S0140673601062523>

760 Salmond, J. A., Tadaki, M., Vardoulakis, S., Arbuthnott, K., Coutts, A., Demuzere, M., Dirks, K. N., Heaviside, C., Lim, S., Macintyre, H.,
761 McInnes, R. N., Wheeler, B. W., 2016. Health and climate related ecosystem services provided by street trees in the urban environment.
762 *Environmental Health* 15(Suppl 1):36.

763 Schelhaas, M., Varis, S., Schuck, A., Nabuurs, G., 2006. Efiscen inventory database. Tech. rep., European Forest Institute, Joensuu, Finland,
764 http://www.efi.int/portal/virtual_library/databases/efiscen/.

765 Seitz, J., Francisco Escobedo, F., 2009. Urban trees and allergies in North Florida. Tech. rep., This document is FOR 206, one of a series of the
766 School of Forest Resources and Conservation Department, UF/IFAS Extension.
767 URL (accessed 24 April 2015) <http://edis.ifas.ufl.edu>

768 Skjøth, C. A., Baker, P., Sadys, M., Adams-Groom, B., 2015a. Pollen from alder (*alnus* sp.), birch (*betula* sp.) and oak (*quercus* sp.) in the {UK}
769 originate from small woodlands. *Urban Climate* 14, Part 3, 414 – 428, new Sensing Technologies and Methods for Air Pollution Monitoring.
770 URL <http://www.sciencedirect.com/science/article/pii/S2212095514000716>

771 Skjøth, C. A., Bilinska, D., Werner, M., Adams-Groom, B., Drzeniecka-Osiadacz, A., Dec. 2015b. Footprint areas of pollen from alder (*Alnus*) and
772 birch (*Betula*) in the UK (Worcester) and Poland (Wroclaw) during 2005-2014. *Acta Agrobotanica* 68, 315–324.

773 Skjøth, C. A., Geels, C., Hvidberg, M., Hertel, O., Brandt, J., Frohn, L. M., Hansen, K. M., Hedegaard, G. B., Christensen, J. H., Moseholm, L.,
774 2008. An inventory of tree species in Europe-An essential data input for air pollution modelling. *Ecological Modelling* 217 (3-4), 292–304.

775 Skjøth, C. A., Ørby, P. V., Becker, T., Geels, C., Schlünssen, V., Sigsgaard, T., Bønløkke, J. H., Sommer, J., Sjøgaard, P., Hertel, O., 2013. Identifying
776 urban sources as cause of elevated grass pollen concentrations using gis and remote sensing. *Biogeosciences* 10 (1), 541–554.
777 URL <http://www.biogeosciences.net/10/541/2013/>

778 Skjøth, C. A., Sikoparija, B., Jager, S., EAN-Network, 2012. Pollen sources. In: Sofiev, M., Bergmann, K.-C. (Eds.), *Allergenic pollen: a review*
779 *of the production, release, distribution and health impacts*. Springer, Ch. 2, pp. 9–27.

780 Skjøth, C. a., Smith, M., Šikoparija, B., Stach, A., Myszkowska, D., Kasprzyk, I., Radišić, P., Stjepanović, B., Hrga, I., Apatini, D., Aug. 2010. A
781 method for producing airborne pollen source inventories: An example of Ambrosia (ragweed) on the Pannonian Plain. *Agricultural and Forest*
782 *Meteorology* 150 (9), 1203–1210.
783 URL <http://linkinghub.elsevier.com/retrieve/pii/S0168192310001292>

784 Skjøth, C. A., Sommer, J., Stach, A., Smith, M., Brandt, J., 2007. The long-range transport of birch (*Betula*) pollen from Poland and Germany
785 causes significant pre-season concentrations in Denmark. *Clinical & Experimental Allergy* 37 (8), 1204 – 1212.

786 Smith, M., Cecchi, L., Skjth, C., Karrer, G., Šikoparija, B., 2013. Common ragweed: a threat to environmental health in Europe. *Environment*
787 *International* 61, 115 – 126.
788 URL <http://www.sciencedirect.com/science/article/pii/S0160412013001682>

789 Smith, M., Emberlin, J., 2005. Constructing a 7-day ahead forecast model for grass pollen at north London, United Kingdom. *Clinical & Experi-*
790 *mental Allergy* 35 (10), 1400–1406.
791 URL <http://dx.doi.org/10.1111/j.1365-2222.2005.02349.x>

792 Smith, M., Emberlin, J., 2006. A 30-day-ahead forecast model for grass pollen in north London, United Kingdom. *International Journal of Biome-*
793 *teorology* 50 (4), 233.
794 URL <http://dx.doi.org/10.1007/s00484-005-0010-y>

795 Sofiev, M., Belmonte, J., Gehrig, R., Izquierdo, R., Smith, M., Dahl, A., Siljamo, P., 2012. Airborne pollen transport. In: Sofiev, M., Bergmann,
796 K.-C. (Eds.), *Allergenic Pollen: A Review of the Production, Release, Distribution and Health Impacts*. Springer, Ch. 5, pp. 127–159.

797 Spot Vegetation, 2017. Spot Vegetation, last accessed 06/03/2017.
798 URL <http://www.vgt.vito.be/pages/VegetationProgramme/generaldescription.htm>

799 Suring, K., Bach, S., Bossmann, K., Wolter, E., Neumann, A., Straff, W., Hflich, C., 2016. PM10 contains particle-bound allergens: dust analysis
800 by flow cytometry. *Environmental Technology & Innovation* 5, 60 – 66.
801 URL <http://www.sciencedirect.com/science/article/pii/S2352186416300049>

802 Thibaudon, M., Šikoparija, B., Oliver, G., Smith, M., Skjøth, C. A., Feb. 2014. Ragweed pollen source inventory for France — The second largest
803 centre of Ambrosia in Europe. *Atmospheric Environment* 83, 62–71.
804 URL <http://www.sciencedirect.com/science/article/pii/S1352231013008108>

805 Vardoulakis, S., Dimitroulopoulou, C., Thornes, J., Lai, K.-M., Taylor, J., Myers, I., Heaviside, C., Mavrogianni, A., Shrubsole, C., Chalabi,
806 Z., Davies, M., Wilkinson, P., 2015. Impact of climate change on the domestic indoor environment and associated health risks in the UK.
807 *Environment International* 85, 299 – 313.
808 URL <http://www.sciencedirect.com/science/article/pii/S0160412015300507>

809 Vardoulakis, S., Heaviside, C., 2012. Health effects of climate change in the UK. Tech. rep., Health Protection Agency.

810 Werner, M., Geels, C., Kryza, M., Skjøth, C. A., 2016. Using a dynamical approach for implementing Ammonia emissions into WRF-Chem over
811 Europe. Springer International Publishing, pp. 345–350.
812 URL http://dx.doi.org/10.1007/978-3-319-24478-5_57

813 Werner, M., Kryza, M., Geels, C., Ellermann, T., Skjøth, C., 2017. Ammonia concentrations over Europe — application of the WRF-Chem model
814 supported with dynamic emission. *Polish Journal of Environmental Studies* in press.
815 URL <https://eprints.worc.ac.uk/id/eprint/5238>

816 Worcester University, 2017. Worcester University pollen calendar, last accessed 06/03/2017.
817 URL <http://www.worcester.ac.uk/pdfs/pollen-calendar.pdf>

818 Yamada, T., Saito, H., Fujieda, S., 2014. Present state of Japanese cedar pollinosis: the national affliction. *Journal of Allergy and Clinical Immunol-*
819 *ogy* 133 (3), 632 – 639.e5.
820 URL <http://www.sciencedirect.com/science/article/pii/S0091674913017119>

821 Zink, K., Pauling, A., Rotach, M. W., Vogel, H., Kaufmann, P., Clot, B., 2013. EMPOL 1.0: a new parameterization of pollen emission in numerical
822 weather prediction models. *Geoscientific Model Development* 6 (6), 1961–1975.
823 URL <http://www.geosci-model-dev.net/6/1961/2013/>

824 Zink, K., Vogel, H., Vogel, B., Magyar, D., Kottmeier, C., 2012. Modeling the dispersion of *Ambrosia artemisiifolia* L. pollen with the model system
825 COSMO-ART. *International Journal of Biometeorology* 56 (4), 669–680.
826 URL <http://dx.doi.org/10.1007/s00484-011-0468-8>

827 Ziska, L., Knowlton, K., Rogers, C., Dalan, D., Tierney, N., Elder, M. A., Filley, W., Shropshire, J., Ford, L. B., Hedberg, C., Fleetwood, P.,
828 Hovanky, K. T., Kavanaugh, T., Fulford, G., Vrtis, R. F., Patz, J. A., Portnoy, J., Coates, F., Bielory, L., Frenz, D., 2011. Recent warming by
829 latitude associated with increased length of ragweed pollen season in central North America. *Proceedings of the National Academy of Sciences*
830 *of the United States of America* 108 (10), 4248–4251.