

Original citation: Oteros, J., Sofiev, M., Smith, Matt, Clot, B., Damialis, A., Prank, M., Werchan, M., Wachter, R., Weber, A., Kutzora, S., Heinze, S., Herr, C.E.W., Menzel, A., Bergmann, K-C., Traidl-Hoffmann, C., Schmidt-Weber, C.B. and Buters, J.T.M. (2019) *Building an Automatic Pollen Monitoring Network (ePIN): Selection of Optimal Sites by Clustering Pollen Stations*. *Science of The Total Environment*, 688. 1263 - 1274. ISSN 0048-9697 (In Press)

Permanent WRaP URL:

<https://eprints.worc.ac.uk/id/eprint/8348>

Copyright and reuse:

The Worcester Research and Publications (WRaP) makes this work available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRaP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

This is an Accepted Manuscript of an article published by Elsevier in *Pharmacological Research*, available online: <https://www.sciencedirect.com/science/article/abs/pii/S1043661819307790>. © 2019 Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International. <http://creativecommons.org/licenses/by-nc-nd/4.0/>

A note on versions:

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRaP URL' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact wrapteam@worc.ac.uk

Building an automatic Pollen Monitoring Network (ePIN): Selection of optimal sites by clustering pollen stations

Jose Oteros¹; Mikhail Sofiev²; Matt Smith³; Bernard Clot⁴; Athanasios Damialis⁵; Marje Prank²; Matthias Werchan⁶; Reinhard Wachter⁶; Alisa Weber⁷; Susanne Kutzora⁷; Stefanie Heinze⁷; Caroline E. W. Herr⁷; Annette Menzel^{8,9}; Karl-Christian Bergmann⁶; Claudia Traidl-Hoffmann^{5,10}; Carsten B. Schmidt-Weber¹; Jeroen T. M. Buters^{1*}

1. Center of Allergy & Environment (ZAUM), Member of the German Center for Lung Research (DZL), Technische Universität München/Helmholtz Center, Munich, Germany.
2. Finnish Meteorological Institute (FMI), Helsinki, Finland.
3. Institute of Science and the Environment, University of Worcester, UK.
4. Federal Office of Meteorology and Climatology MeteoSwiss, Payerne, Switzerland.
5. Chair and Institute of Environmental Medicine, UNIKA-T, Technical University of Munich and Helmholtz Zentrum M., Augsburg, Germany.
6. Foundation German Pollen Information Service (PID), Berlin, Germany.
7. Bayerisches Landesamt für Gesundheit und Lebensmittelsicherheit (LGL), Munich, Germany.
8. Technische Universität München, Ecoclimatology, Department of Ecology and Ecosystem Management, Freising, Germany.
9. Technische Universität München, Institute for Advanced Study, Garching, Germany.
10. Christine Kühne Center for Allergy Research and Education (CK Care), Davos, Switzerland.

* Corresponding author:

Jeroen Buters

ZAUM-Technical University Munich

Biedersteinerstrasse 29

80802 Munich, Germany

Buters@tum.de

+49-89-41403487

Abstract

Airborne pollen is a recognized biological indicator and its monitoring has multiple uses such as providing a tool for allergy diagnosis and prevention. There is a knowledge gap related to the distribution of pollen traps needed to achieve representative biomonitoring in a region. The aim of this manuscript is to suggest a method for setting up a pollen network (monitoring method, monitoring conditions, number and location of samplers etc.). As a case study, we describe the distribution of pollen across Bavaria and the design of the Bavarian pollen monitoring network (ePIN), the first operational automatic pollen network worldwide.

We established and ran a dense pollen monitoring network of 27 manual Hirst-type pollen traps across Bavaria, Germany, during 2015. Hierarchical cluster analysis of the data was then performed to select the locations for the sites of the final pollen monitoring network. According to our method, Bavaria can be clustered into three large pollen regions with eight zones. Within each zone, pollen diversity and distribution among different locations does not vary significantly. Based on the pollen zones, we opted to place one automatic monitoring station per zone resulting in the ePIN network, serving 13 million inhabitants. The described method defines stations representative for an homogeneous aeropalynologically region, which reduces redundancy within the network and subsequent costs (in the study case from 27 to 8 locations). Following this method, resources in pollen monitoring networks can be optimized and allergic citizens can then be informed in a timely and

effective way, even in larger geographical areas.

Keywords: Aerobiology; air quality; automatic pollen monitoring; BAA500; biomonitoring network; pollen

1. Introduction

Pollen is part of the biological exposome, carrying allergens, fungi and bacteria able to activate the human immune system (Buters et al., 2015; Buters et al., 2012; Galán et al., 2013; Oteros et al., 2019). There are a number of networks that routinely monitor airborne pollen worldwide. These were built for a range of purposes (Buters, 2014), such as examining gene flow (Hofmann et al., 2014), allergy prevention (de Weger et al., 2014; Fernandez-Rodriguez et al., 2014; Sofiev et al., 2017; Werchan et al., 2017; Zink et al., 2012), crop forecasting, pest control, impacts of land use changes (Aguilera and Ruiz-Valenzuela, 2014; Cunha et al., 2016; Dhiab et al., 2016; Jochner-Oette et al., 2017; Rodríguez-Rajo et al., 2010), climate change impacts (De Linares et al., 2017; Galán et al., 2016; Smith et al., 2014; Zhang et al., 2015; Ziello et al., 2012a; Ziello et al., 2012b) and monitoring biodiversity (Belmonte et al., 2000; Cariñanos et al., 2013; Sikoparija et al., 2009; Thibaudon et al., 2014).

The simplest way to sample airborne pollen and fungal spores is to collect the particles deposited or impacted on certain surfaces (Darwin, 1846). First generation traps (1G) do not provide volumetric information (Cour, 1974; Durham, 1946; Ogden and Raynor, 1967), although some have been termed semi-volumetric because they were calibrated with help of an anemometer (Orlandi et al., 2014). On the other hand, second generation samplers (2G) use a range of different sampling principles (Mandrioli et al., 1998) and provide volumetric data making them comparable. The system designed by Hirst more than 65 years ago is based on the impaction principle (Hirst, 1952), and is the standard in pollen and fungal spore monitoring networks in most of the world (Galán et al., 2014; VDI4252-4, 2016).

The Hirst-type trap has several advantages, e.g. it delivers pollen concentration data with a temporal resolution of up to 2 h, it provides volumetric information (i.e. pollen/m³) and it has a long autonomy of up to 7 days.

There are currently more than 600 Hirst-type traps actively running worldwide, mostly in Europe (<https://www.zaum-online.de/pollen/pollen-monitoring-map-of-the-world.html>); (Buters J.T.M. et al., 2018). A standardized sampling system and working methodology is essential for a network, and there are numerous publications on the standardization of this monitoring method (Alcázar et al., 1999; Comtois et al., 1999; Galán and Domínguez-Vilches, 1997; Gharbi et al., 2017; Levetin et al., 2000; Maya-Manzano et al., 2017; Oteros et al., 2017; Oteros et al., 2013a; Sikoparija et al., 2016; Smith et al., 2019; VDI4252-4, 2016). The “Minimum requirements for aerobiology” (Galán et al., 2014), supported by the European Aerobiology Society, is recognized as an international standard for pollen monitoring using Hirst-type traps. Other devices currently used in active networks include the Cour method (Oteros et al., 2014), Rotorod samplers (Portnoy et al., 2004) and Durham traps (Teranishi et al., 2000).

The main disadvantage of all manual methods is that data production is highly time consuming and thus it is not feasible to provide timely information with respect to the current, real-life situation. A time lag of several days between actual pollen flight and reported pollen flight is common. A solution could be automation, but the complexity of pollen identification has made this impossible until now. New technologies allow for fully automatic-online pollen monitoring with third generation automatic traps (3G). The main advantages of these traps are that they can provide almost real-time information and the data are free of random errors produced by human interferences. Such 3G automatic systems are starting to be used for routine pollen monitoring, e.g. the KH-3000 (Kawashima et al., 2017), the Plair PA-300/Rapid E (Crouzy et al., 2016; Šaulienė et al., 2019), Pollen Sense (<http://pollensense.com/>), the BAA500 (Oteros et al., 2015b) or WIBS-4 (O’Connor et al., 2014).

The main reason for building the Bavarian pollen monitoring network ePIN was to reduce delays in the dissemination of pollen information and to inform allergic citizens, physicians, and health organizations in a timely manner. Another aim of the ePIN network was to serve as a registry of environmental changes. To do so, we had to increase the representativeness of pollen monitoring across Bavaria. The whole federal state has actually a population of 13 million inhabitants, but airborne pollen has routinely been monitored at only three locations (www-genesis.destatis.de). A handicap of classical pollen monitoring networks is that they are often built on the basis of stations set up by individuals, a fact that does not always guarantee an appropriate choice of positions for pollen monitoring stations. There is a general lack of studies on the optimization of monitoring locations, and so the decision where to install a station has usually been based on personal preferences, often in large urban areas (Buters et al., 2018).

The aim of this study was to the overall “Aims” were to provide near real-time health information to pollen allergic individuals and health care practitioners and so serve as a valuable source of biological data for (but not limited to) long-term biodiversity and climate change studies.

The following are “Objectives” or steps taken to achieve the overall aims:

- Design a method for building a pollen network considering monitoring method, homogeneous monitoring conditions, number and location of monitoring stations.
- Investigate the pollen distribution across Bavaria.
- Use this information to produce the first operational automatic pollen network in history: the electronic Pollen Information Network (ePIN).

2. Material and Methods

2.1 Study area

Bavaria extends to an area of 70.553 km² being the largest federal state in Germany, and is located in the south of the country (Figure S1). The maximal distance between locations in Bavaria is more than 400 km. The environmental zones are quite heterogeneous, containing a wide range of climates from the cold Bavarian Alps in the South (Zugspitze with an annual mean temperature of -4.3°C and mean precipitation of 2071 mm) to the warmer-dryer Franconian wine area in the North-West (Würzburg with an annual mean temperature of 9.6°C and mean precipitation of 601 mm). The annual mean temperature at Munich is 8.7°C with a mean annual precipitation of 834 mm and the annual mean temperature at Nuremberg is 9.3°C and mean precipitation is 637 mm (all climate data from the German Meteorological Service for the reference period 1981-2010). These different conditions render representative pollen monitoring across the state rather complex.

Thirty-seven percent of the Bavarian area is covered by forests (<https://bwi.info/>). The most important forested areas are the Alps (South) and the Bavarian Forest (West), both with coniferous predominance. The most predominant pollen types in the region were from *Pinus* and *Picea*. Also, pollen from deciduous trees in forest patches were dispersed across the state, such as *Betula* and *Fagus*. The Bavarian environment is also characterized by extensive pastures with an abundance of anemophilous herbaceous plants as from the families Poaceae and Plantaginaceae. The pollination period in Bavaria is expected to occur from the early flowering anemophilous taxa such as *Alnus* and Cupressaceae in January to the latest flowering anemophilous genera of *Ambrosia* in October.

2.2 Proposed steps of building the network

2.2.1 Selection of pollen monitoring methods

We reviewed the advantages and disadvantages of the current options for pollen monitoring (see results). For the final (permanent) ePIN network, an automatic pollen monitoring system was selected, whereas for the pilot (test) network we chose the Hirst-type monitoring method (Hirst, 1952), as it is accepted currently as the most standard pollen monitoring system.

2.2.2 Review on the site conditions for installing a pollen monitoring device

We performed a bibliographical review and an international survey among the administrators of pollen monitoring networks about the main factors affecting the decision of trap location at the local (site) scale (see results).

2.2.3 Building a redundant pollen network (pilot network)

To select the final number and location of monitoring stations across Bavaria, we first built a dense network of 27 Hirst traps throughout the country (the pilot network) and, based on the results obtained, reduced the number of locations with hierarchical clustering (Ward's clustering method). This mathematical method has already been used to set up chemical air quality monitoring networks (Nakamori and Sawaragi, 1984). The number of stations was deliberately selected to configure a redundant network that could be collapsed after the analysis. To ensure it was a redundant network, the density of stations was the highest in the world for a pollen network of these dimensions (27 stations in Bavaria - 70,550.19 km²) (Buters et al., 2018).

2.2.3.1 Selection of monitoring locations in the pilot network

The pre-selection of 27 pollen monitoring locations was performed in order to increase the regional representativeness of the pollen sampling. The following factors were considered: demography, availability of historical time series, climate, land use types, topography and proximity to local pollen sources. The pre-selection was done trying to satisfy most of the specific features that the network should have:

- Must preserve historical time series, so historical locations were included.
- Must be informative for the majority of the population, so the most populated areas were closely monitored. Bavaria is unevenly populated, with several major urban agglomerations (Figure S2a). The 25 km radius buffer around all the pre-selected stations covered 94% of the Bavarian population. It is accepted that the area influencing a roof-level trap is >25 km (Oteros et al., 2015a).
- Must be capable of detecting pollen episodes early, so source areas and borders of the state were preferred (Figure S2b). The places near main pollen sources are more appropriate locations to perform a better forecasting of airborne particle distribution. Broad-leaved forests consisting of early spring flowering allergenic trees, like alder and birch, are present in patches, and are spread over Bavaria and near its borders (i.e. the locations were located close to the broad-leaved forest patches and the big forest areas as the North-West broad-leaved forest, the Bavarian forest and the Alps). This suggests the possibility of setting stations closer to the borders of Bavaria to earlier catch the moments of the forests starting to flowering out of the boundaries. Grasses are also common in Bavaria; their pollen is known to be less efficiently transported than those from anemophilous trees.
- Must provide data for model-based forecasting, for data assimilation and for the model evaluation.
- Must cover the major biogeographic environments existing in Bavaria. Temperature is important

for the season start and duration: plants in the warmer parts tend to flower earlier. In this sense, the warmest regions of Bavaria are north-west and south-east. The colder and wetter areas are the alpine mountains in the south and the area of the Bavarian Forest in the east (Figure S2c, S2d).

The pre-selection of the locations for new stations was carried out for zones with a 25 km radius (this number is smaller than the accepted influence area of a pollen trap at roof level (Oteros et al., 2015a)). The selection of the specific location was then done by screening for homogeneous monitoring conditions inside each pre-selected zone within the network. To ensure a proper coverage of all pollen sources in Bavaria, an analysis with the System for Integrated modeLLing of Atmospheric coMposition (SILAM, <http://silam.fmi.fi>) was conducted, ensuring a large coverage of Bavarian sources (Sofiev et al., 2013; Sofiev et al., 2015). The footprint that the pilot network covered in 2015 is shown in Figure 1.

A footprint is the area comprising the sources that affected the monitored parameter during a specific observation period (in the current case, pollen data of every specific day during the 2015 season). Such a footprint delineates the area where the sources would contribute to the monitor readings, if emitting pollen during the corresponding time. Areas outside the footprint do not affect the pollen monitor. This “negative” part of the footprint message makes it a handy tool for delineating the network weaknesses. Formally, the convolution of emission E , and the footprint “intensity” φ^* over space and time is equal to the mean concentration C at the specific place during the specific period (monitor location and time of activity) (function 1).

$$\int_T \iiint \varphi^* E d\mathbf{r} dt = C$$

(1)

Thus, a footprint delineates the area where the sources would contribute to the monitor

readings if emitting pollen during the corresponding time and shows which sources can affect the monitor during the specific observation period. The sum of the footprints of all daily measurements by the 27 ePIN stations in 2015 is shown in figure 1. The footprints were calculated over three days backwards from the observation day, i.e. covered the areas from where the emitted pollen needed up to three days to reach the monitor.

>> **Figure 1**

The exact location to set up a pollen trap was selected by a site visit of potential places within the 25km-radius zones. Criteria for a site were reviewed in step 2.2.3.2 and are listed in Table 2. This resulted in 27 locations described in Table 1.

>> **Table 1**

2.2.3.2 Establishment and quality control in the dense pilot pollen monitoring network

The network of 27 Hirst pollen traps was set up across Bavaria at the selected locations (Table 1) during the winter of 2014-2015. The network then operated from the end of February until the end of September 2015. All traps were located so that homogeneous monitoring conditions may be achieved (see Table 2). For instance, all stations were built at 12 m a.g.l. (± 3 m) eliminating the large variability of the first 10 m layer and eliminating differences between stations in height (Rojo et al., 2019a), all traps were also located at 1.5 m above roof level by a standard tower (Figure S3a) and at least 2 m from the building edge. Flow rates of the pollen traps were calibrated using the same flowmeter thereby reducing intra-rotameter variability (Oteros et al., 2017). Drums and microscope slides were processed centrally by a single laboratory (figure S3b) under identical conditions. The drums were sent bi-weekly to the 27 monitoring stations. Slides were processed using the standard operating procedure described by Galán et al. (Galán et al., 2007).

Each pollen slide corresponded to one independent day (Figure S3c). All slides had a blue line, marking midday (12:00). Four blue dots were set 1mm apart from each other at the center of the slide with a standard self-designed tool to guide the analyst for the starting point of each horizontal line. Pollen microscopic identification and counting were conducted in four continuous horizontal sweeps along the whole slide under 400x magnification, each sweep starting from each one of the marked points.

Sub-sampling of the slide is essential for reducing workload. The area of the slide sub-sampled during the analysis was at least 7% of the slide, following the recommendations of VDI guidelines (VDI4252-4, 2016). The use of a standard 12.5 mm net micrometer reduced the area of the slide examined to 9% when examining 4 transects (or 7% when using a 10mm net micrometer) (figure S2d). A standard correction factor was used to reduce error. Pollen counts (raw data) were entered into a specially designed computer program to reduce typing errors (figure S2f). Data were exported from this program and stored in an online SQL ZAUM database.

In total, 13 pollen types were analyzed: *Alnus*, *Ambrosia*, *Artemisia*, *Betula*, *Carpinus*, *Cupressaceae*, *Fraxinus*, *Picea*, *Pinus*, *Plantago*, *Poaceae*, *Populus*, and *Urticaceae*. Pollen were reported in 12-hour concentrations for each station during the study period. Pollen not falling within the 13 specified pollen types were reported as “unspecified” pollen grains.

An external Quality Control program of the analysts was performed with a novel method, published in detail by Smith et al. (Smith et al., 2019).

2.2.4 Establishment of the final pollen monitoring network

2.2.4.1 Selection of the definitive number of traps by clustering analysis

We clustered the information obtained from all pollen traps in order to determine areas with a similar distribution in pollen. Due to the complexity of data (daily pollen monitoring of 27 stations and 13 pollen types), we applied a multivariate statistical method able to consider all the variables at the same time.

First, we preselected the part of the database to be included in the analysis. We selected the most abundant pollen types for the clustering analysis (>1000 pollen grains/season on average): *Betula*, Cupressaceae, *Fraxinus*, *Pinus*, *Picea*, Poaceae and Urticaceae. Those are the pollen types with the largest spread of annual pollen values, whereas less abundant pollen types do not show big differences among the locations and so it makes little sense to cluster pollen zones based on them (Figure 2A). The flowering dates by pollen type is shown in Figure 2B, carried out with the R package AeRobiology (Rojo et al. 2019). From the 27 monitoring stations, DEERLA was excluded from the analysis due to technical malfunctions. Cupressaceae pollen was excluded from the analysis due to the incomplete monitoring of the whole season. *Pinus* was excluded from the clustering analysis to avoid overrepresentation of the Pinaceae family for designing the network, *Picea* was included. The pollen season for each pollen taxon was defined for the whole Bavaria as follows, excluding the long tails before and after the season with zero values: *Betula* (from 1/4 to 14/5); *Fraxinus* (from 25/3 to 2/5); *Picea* (from 22/4 to 26/5); Poaceae (from 8/5 to 6/8); Urticaceae (from 1/6 to 8/9).

>> Figure 2

Second, we calculated all Pearson correlations in daily pollen concentrations between pairs of stations (26 stations). We applied this correlation analysis for each selected pollen type (5 pollen types in total: *Betula*, *Fraxinus*, *Picea*, Poaceae and Urticaceae). For each pollen taxon, we only included in the

analysis the stations with at least 80% of the data during the season. When a correlation was too low, as could happen by chance, all correlations <0.5 were equalized to 0 in the correlation matrixes for the clustering analysis.

Third, we applied a clustering analysis to the correlations' coefficients (Hierarchical clustering by Ward's method) (Oteros et al., 2013b; Ward Jr and Hook, 1963). In the analysis, 26 cases were included to be conglomerated (each monitoring station). Each variable was defined as the correlation coefficient between one station (for each pollen type) and each one of the 26 stations (for the same pollen type). Twenty-six cases were included in the analysis (one per station). A correlation coefficient is not a metric of distance *per se*, but the combination of them allows us to calculate Euclidean distances. Furthermore, a visualization of the five closest Euclidean distances for each element is shown by a network plot (see figure 4).

2.2.4.2 Determination of the final monitoring locations

Within each cluster calculated in step 2.2.4.1, one station was then selected. We selected the most relevant station of each cluster using the following selection criteria:

- The station covering the highest **population** was selected (If two or more stations differ by **$<0.5\%$ population**, they are all selected at this stage).
- In the case of a draw (similar population), the station **closest to the border of Bavaria** was selected.
- Two selected stations cannot be located closer than **70 km apart** (ensuring a proper coverage of the whole surface). If two stations are closer than 70km, then the most populated location is selected in one sub-cluster and the next by population is selected in the other cluster.

- Four stations using the Hirst-type biomonitoring method were kept as a parallel manual network to maintain a historical time series (DEOBER, DEMUST, DEBAMB and DEUFS) and were not selected for the automatic network.

An automatic network was then built in Bavaria based on these criteria.

3. Results and Discussion

3.1 Selection of the pollen monitoring method

The main goals of this pollen monitoring network were: 1. to provide near real-time health information to pollen allergic individuals and health care practitioners. 2. To serve as a valuable source of biological data for (but not limited to) long-term biodiversity and climate change studies. Hence, the network must allow for, if possible, the continuation of historical time series and monitor the whole spectrum of airborne pollen types, not only the allergenic ones.

A permanent network of Hirst-type traps may provide pollen information with a minimum lag time of 1 to 2 days, involving a colossal human effort and huge costs (arising mostly from the need of experienced personnel). We therefore focused on an alternative, automated, detection system able to provide near-real-time information. There is a well-established monitoring network of 120 traps based on an automatic-online monitoring system in Japan, the KH-3000 (Kawashima et al., 2017), but this automatic system until now has been unable to provide accurate information on the complete range of pollen diversity. Other systems look promising for the development of a complete and reliable automatic recognition system of pollen: e.g. Pollen Sense (<http://pollensense.com/>, [no scientific publication available](#)), BAA500 (Oteros et al., 2015b), Plair PA-300 Rapid E (Crouzy et al., 2016; Šaulienė et al., 2019) or Swisens (www.swisens.ch, [no scientific publication available](#)). Of these, only two automatic pollen monitoring systems were fully operational on the date of the network setup, BAA500 (Oteros et al., 2015b) and Plair PA-300 (Crouzy et al., 2016).

The BAA500 was the preferred monitoring system for the ePIN network because it has specific features that make it a good candidate for the transition from manual to automatic monitoring. The BAA500 uses image recognition emulating the process of a human using a microscope, and can provide

3-hour pollen concentrations online (it can be programmed to deliver 1 hour concentrations, but not checked the error to date) and its averaged identification error rate is below 10% (Oteros et al., 2015b).

3.2 Selection of homogeneous pollen monitoring conditions

The results of the bibliographical review and the survey among European experts about suggested conditions for pollen monitoring are shown in Table 2. Most of these criteria are extracted from the pre-established knowledge about pollen monitoring (Galán et al., 2007; Galán et al., 2014; Mandrioli et al., 1998; Saar and Meltsov, 2011).

>> Table 2

Some criteria were not defined quantitatively because they have not been exhaustively studied. In this sense, the criteria of Table 2 should not be termed as optimal but more motivated by the necessity of defining comparable standard conditions for the whole network.

3.3 Running the dense pilot pollen network

At a regional scale, the pollen stations had to be placed in areas of interest to the general population, thus demography was a main criterion. Other factors that strongly affect air quality such as topography (Rojo and Perez-Badia, 2014), land use (Haberle et al., 2014) or weather (Damialis et al., 2005), which modify the interpretation of pollen monitoring, were also considered.

The pilot pollen monitoring network based on these criteria is shown in Figure 3. Based on these criteria, the steps followed to perform the selection were:

- Existence of historical time series was considered for monitoring climate change impacts. **9 Stations** with historical time series were selected at first: DEZUSM, DEMUST, DEBAMB, DEERLA, DEBAYR, DEDONA, DEMUNC, DEGAIS and DEBERC.

- Existence of other independent stations running during 2015 in Bavaria were also considered as part of the pilot network. **5 Stations were added:** DEBIED, DEAUUGS, DEPFRO, DEGARM and DEUFS.
- The network must be capable of detecting pollen episodes early, so source areas got particular attention. The places closer to the main pollen sources are more appropriate locations to perform a better forecasting of airborne particles. **4 Stations were added:** DEALTO (Close to the Austrian sources), DEVIEC (Bavarian Forest), DEMARK (North-West Franconian forest) and DEOETT (Schwaben-South Franconia forest).
- The network must be informative for the bulk of the population, so the most populated areas were closely monitored when necessary. **4 Stations were added:** DEFEUC (Close to Nürnberg), DEKITZ (Close to Würzburg), DEKOES (Close to Ingolstadt) and DEMIND (populated area in Schwaben).
- Must cover the major biogeographic environments existing in Bavaria. Most of the environments were already covered. **5 Stations were added:** DELAND (middle-stream Danube area), DETROS (South East Bavaria), DEPASS (downstream Danube area), DEWEID and DEHOF (North-East Bavaria).

Out of those stations, five were independently managed: DEAUUGS (UNIKA-T, first data: 1999), DEBIED (private-ZAUM, first data: 2003), DEOBER (private-PID, first data: 1982), DEGARM (TUM, first data: 2008) and DEUFS (TUM, first data: 2008). Two stations were managed by PID with long time series: DEMUST (first data: 1990) and DEMUNC (first data: 1987). The other 20 stations were managed at the central lab of ZAUM (these raw data are supported with the manuscript). From these 20 stations, 7 correspond to historical PID locations with long time series but previously discontinued: DEBAMB (first data: 1989), DEBAYR (first data: 1989), DEERLA (first data: 1987), DEDONA (first data: 1989), DEZUSM (first data: 1987), DEGAIS (first data: 1992) and DEBERC (first data: 1987). Thirteen stations corresponding to new locations commenced in 2015: DEHOF, DEMARK, DEKITZ, DEWEID, DEFEUC, DEOETT, DEVIEC, DEKOES, DELANDS, DEPASS, DEALTO, DETROS and DEMIND.

>> **Figure 3**

3.4 Clustering pollen zones and selected locations of the definitive ePIN network

The result of the analysis is summarized as a dendrogram (Fig. 4a), starting with 26 elements and ending with one (senseless) big cluster. We had to determine the number of clusters that represented similarities between locations and clustering distance. Elbow plot suggest the existence of 4 main clusters. After 4 clusters, the variance explained by additional clusters is smaller (Figure S4).

Figure 4a shows four clearly defined clusters: A Central Cluster (C1) with 12 stations distributed in the center of Bavaria; A Cold Cluster (C2) with 6 stations distributed in the colder areas (Alps and Bavarian Forest); An Outlier Cluster (C3) with 3 stations without apparent connection; A Franconian Cluster (C4) with 5 stations distributed in the North. The abovementioned names of each cluster are provisional and serve only for providing a better understanding of the grouping.

>> **Figure 4**

There were not always well-defined boundaries between clusters and, so, some regions could be considered as transitional or subclusters (8) inside the bigger four clusters. For instance, in the cluster C4 (Franconian cluster), DEHOF and DEBAYR are in a small sub-group (East-Franconia) because of bioclimatical similarities (lower temperatures, see Figure S3c) with respect to West-Franconia. In the same way, DEBIED and DEMUNC are the nearest stations in the first cluster C1 (Central cluster) and indeed both traps are located within the same city (Munich). DEALTO, DEPASS and DETROS showed similar data, the three locations are quite close to the southeast of Bavaria. DEAUGS and DEMIND showed similar data too, both locations are close in southwest Bavaria. DEGARM and DEUFS constitute also a small subcluster inside the C2 (Cold cluster), both locations being only 10 km apart in the Alps region, however altitudinally differing by 1900 m. Three stations, DEMUST, DEZUSM and DELANDS (Cluster C3) were considered as outliers (i.e. the cluster was formed of stations located in distant and

different bioclimatic areas).

Figure 4b shows a visualization of the Euclidean distances between locations (based on all the pairs of correlations), for a better understanding of sub-clusters and transitional areas. For visualization, we represented only the edges with the five closed elements. Figure 4b is a way of visualizing a multidimensional space into two dimensions by rescaling the weight of the edges, so the result does not necessarily match with the dendrogram produced by Hierarchical clustering. This visualization allows us to understand that inside C4, the stations of East-Franconia (DEHOF and DEBAYR, colder area) are closer to the Cold cluster (C2) and the stations of the warmer area are closer to the Central cluster (C1). Inside the cluster C2, DEVIEC is closer to the central cluster, indeed this station is at the border of C1 surrounded by DEWEID, DEDONA and DEPASS-DEALTO-DETROS. As can be observed, DEUFS is the station farthest away from the rest of the network inside C2, as this station is located under extreme conditions at the top of the Alps at 2656 m a.s.l., being the highest pollen monitoring station in the world (<http://www.schneefernerhaus.de/startseite.html>). The three stations included into the Outlier cluster (C3) appear isolated also in Figure 4b. DEWEID appears connected with the colder stations of C4 and the 5 stations of C2, and indeed this station is also located in the Bavarian Forest, under transitional conditions between C1 and the stations DEHOF-DEBAYR (C4) and DEVIEC (C2). The hierarchical clustering put this station in a subcluster together with DEFEUC (both are the northern stations of C1).

For each of the 8 subclusters we selected only one station for automation in the permanent network. To select the most relevant station inside each sub-cluster we followed a series of selection criteria. Table 3 shows the final selection.

>> Table 3

The station DEOBER would have been selected for the permanent network by population, however this station has one of the longest time series in Bavaria (Simoleit et al., 2016), thus the station

was already been selected for a parallel permanent manual network. To avoid double sampling at the same location and to save resources, we defined the fourth criteria, the station was changed for the next suitable station inside the alpine sub-cluster according to the selection criteria (DEGARM). This station has a special touristic-economic interest in a changing environment (Hamilton and Tol, 2007). Figure 5 shows the final selected network with 8 automatic locations.

>> Figure 5

During the last 60 years of pollen monitoring, the standardization of methods was an important issue and all the efforts ended into a high degree of comparability between pollen data across the globe (Galán et al., 2014). At the same time, there was no evolution in the sampling technology, predominantly using the same pollen sampler (i.e. the Hirst-type trap) with the same features as the original design (Oteros et al., 2017). The building of a network based on an automatic system is an alternative and promising option. First, it provides information about airborne pollen in almost real time, eliminating the workload and the delay of the information, which are the main disadvantages of classical pollen analysis for health purposes. It also eliminates human variability and personal bias during routine monitoring, increasing the comparability of the data. Different 3G automatic systems will probably coexist during the following decades without any becoming dominant. Pollen experts will be essential to calibrate, supervise and support machines to be adapted to changing environments. In our vision, classical pollen monitoring will not disappear, but will be performed more selectively by pollen experts and only for specific scientific purposes. Although the automatic data flow is already creating new problems, like the necessity of filtering the disseminated information (Bastl et al., 2017), the advantages provided by the automatic monitoring was always an ambition of aerobiologists, now becoming possible.

We established a method to determine how many traps are needed to represent the pollen distribution within a certain area. Our method minimizes effort and operational costs whilst providing a

representative picture of the pollen flight within an area. Of course, we would be able to improve monitoring of pollen flight if we had unlimited budget. Furthermore, by automating the monitoring method we were able to obtain more rapid data delivery and provide better service to allergic individuals.

4. Conclusions

- The first operational automatic pollen network in the world was built in Bavaria (ePIN network), based on the automatic system BAA500.
- Collapsing a dense network by Ward's clustering analysis, the minimal number and position of monitoring locations were assessed.
- Standard conditions for pollen monitoring were reviewed and summarized.
- Bavaria (Germany) can be clustered in 3 pollen zones and 8 sub-zones regarding pollen distribution and abundance.
- In the studied pilot network, the most abundant pollen types in Bavaria were those from *Pinus*, *Betula*, *Urticaceae* and *Poaceae*.
- In the studied pilot network, the main pollination period in Bavaria ranged from February (*Alnus*) to October (*Ambrosia*).
- In the studied pilot network, the pollen taxa showing the longest pollination period were the herbaceous families of *Poaceae*, *Plantaginaceae* and *Urticaceae*.
- In building a network first the locations and micro-environment of the monitoring stations must be determined, independent of the choice of the instrument to be used.

Acknowledgments

To the ZAUM team who helped building and running the network: Christine Weil, Ramona Hegedüs, Elke Bartusel, Cordula Ebner v. Eschenbach, Joana Candeias, Azahara Martínez, Danijel Kupresanin, Daniel Fritz, Herr Ertl and Gudrum Pusch. We also thank the international advisors and collaborators: Michelle Thibaudon, Roberto Albertini, Uwe Berger, Marco Fornaciari, Fabio Orlandi, Branko Sikoparija, Sevcan Celenk, Letty de Weger, Łukasz Grewling, Frieder Hofmann, Reinhard Wachter, Barbora Werchan and Carmen Galán. To the Bavarian Health and Food Safety Authority– LGL ePIN team: Alisa Weber, Susanne Kutzora, Stefanie Heinze, Tuan-Anh Vu & Caroline Herr. Special acknowledgement to the pollen analysts: Christine Weil, Athanasios Damialis, Reinhard Wachter, M^a José Velasco-Jiménez, Moisés Martínez-Bracero, Mirjana Špehar, Tatjana Pejak-Sikoparija, Anna Schubert, Paula Castiñeiras-Campos, Orsolya Udvardy, Cristina Orihuela-Gómez, Natasa Camprag-Sabo, José M^a Maya-Manzano, Łukasz Kostecki, Weronica Haratym, Malgorzata Nowak, Gloria Florit-Bosch, Gergely Mányoki, Fátima Aguilera-Padilla, Estefanía Sánchez-Reyes, Dorota Myszkowska, Dina Polašek-Hranilović, Estrella Alfaro-Saiz, Agata Szymanska, Deutscher Polleninformationsdienst (PID) pollen analysts. We are very grateful to the responsables of each monitoring location performing the maintenance and management of Hirst traps and/or institutional/scientific support: Jürgen Dietrich, Robert Moser, Alfred Schimmer, Günther Sperber, Volker Melichar, Karl Gerhard, Uwe Pfeiffle, Günter Reinke, Ralf Göller, Joachim Haun, Heinz Ott, Thomas Kurz, Frank Scharnowski-Fischer, Albert Dirnberger, Stefan Nowack, Franz Huber, Joachim Merkel, Roland Buchheit, Michael Pfeifer, Carl-Peter Bauer, Axel Holstege, Stefan Frank, Herr Rau, Herr Markus de Ridder, Herr Kukula, Herr Grassl, Herr Bauer, Leitung Haustechnik, Josef Dietrich, Herr Zeinz, Herr Zapke, Herr Treffer, Herr Tölle, Herr Staudigl, Herr Schuler, Herr Schmauß, Herr Rohreigner, Herr Reitschuster, Herr Preiss, Herr Penka, Herr Kuppler, Herr Köppel, Herr Hümmer, Herr Hinterbrandner, Herr Heiß, Herr Hegwein, Herr Hauer, Herr Hain, Herr Götz, Herr Dormaier, Herr Bittel, Herr Beer, Frau

Wolf, Frau Seidel, Herr Tellkamp. We acknowledge the implication and support from the Ministries: Dr. Martin Hicke, Dr. Johannes Kiesel, Dr. Tanja Gschlößl.

Funding Sources

This study was funded by the Bavarian State Ministry of Health and Care grant G47-G8180.1-2014/5-25 and the Bavarian State Ministry of the Environment and Consumer Protection grant TKP01KPB-69312.

Declaration of competing financial interests (CFI)

The authors declare they have no actual or potential competing financial interests.

References

- Aguilera F, Ruiz-Valenzuela L. Forecasting olive crop yields based on long-term aerobiological data series and bioclimatic conditions for the southern Iberian Peninsula. *Spanish Journal of Agricultural Research* 2014; 12: 215-224.
- Alcázar P, Galán C, Carinanos P, Domínguez-Vilches E. Effects of sampling height and climatic conditions in aerobiological studies. *Journal of Investigational Allergology and Clinical Immunology* 1999; 9: 253-61.
- Bastl K, Berger M, Bergmann KC, Kmenta M, Berger U. The medical and scientific responsibility of pollen information services. *Wien Klin Wochenschr* 2017; 129: 70-74.
- Belmonte J, Vendrell M, Roure JM, Vidal J, Botey J, Cadahía À. Levels of Ambrosia pollen in the atmospheric spectra of Catalan aerobiological stations. *Aerobiologia* 2000; 16: 93-99.
- Buters J. Pollen allergens and geographical factors. In: Akdis C, Agache I, editors. *Global Atlas of Allergy*. 1. European Academy of Allergy and Clinical Immunology (EAACI), Zurich, 2014, pp. 36-37.
- Buters J, Prank M, Sofiev M, Pusch G, Albertini R, Annesi-Maesano I, et al. Variation of the group 5 grass pollen allergen content of airborne pollen in relation to geographic location and time in season. *J Allergy Clin Immunol* 2015; 136: 87-95 e6.
- Buters JTM, Antunes C, Galveias A, Bergmann KC, Thibaudon M, Galan C, et al. Pollen and spore monitoring in the world. *Clinical and Translational Allergy* 2018; 8: 9.
- Buters JTM, Thibaudon M, Smith M, Kennedy R, Rantio-Lehtimäki A, Albertini R, et al. Release of Bet v 1 from birch pollen from 5 European countries. Results from the HIALINE study. *Atmospheric Environment* 2012; 55: 496-505.
- Cariñanos P, de la Guardia CD, Algarra JA, De Linares C, Irurita JM. The pollen counts as bioindicator of meteorological trends and tool for assessing the status of endangered species: the case of *Artemisia* in Sierra Nevada (Spain). *Climatic Change* 2013; 119: 799-813.
- Comtois P, Alcazar P, Néron D. Pollen counts statistics and its relevance to precision. *Aerobiologia* 1999; 15: 19-28.
- Cour P. Nouvelles techniques de detection des flux et des retombées polliniques: Etude de la sedimentation des pollens et des spores a la surface du sol. *Pollen et spores* 1974.
- Crouzy B, Stella M, Konzelmann T, Calpini B, Clot B. All-optical automatic pollen identification: Towards an operational system. *Atmospheric Environment* 2016; 140: 202-212.
- Cunha M, Ribeiro H, Abreu I. Pollen-based predictive modelling of wine production: application to an arid region. *European Journal of Agronomy* 2016; 73: 42-54.
- Damialis A, Fotiou C, Halley JM, Vokou D. Effects of environmental factors on pollen production in anemophilous woody species. *Trees* 2010; 25: 253-264.
- Damialis A, Gioulekas D, Lazopoulou C, Balafoutis C, Vokou D. Transport of airborne pollen into the city of Thessaloniki: the effects of wind direction, speed and persistence. *International Journal of Biometeorology* 2005; 49: 139-45.
- Darwin C. An account of the Fine Dust which often falls on Vessels in the Atlantic Ocean. *Quarterly Journal of the Geological Society* 1846; 2: 26-30.
- De Linares C, Delgado R, Aira MJ, Alcázar P, Alonso-Pérez S, Boi M, et al. Changes in the Mediterranean pine forest: pollination patterns and annual trends of airborne pollen. *Aerobiologia* 2017; 33: 375-391.
- de Weger LA, Beerthuizen T, Hiemstra PS, Sont JK. Development and validation of a 5-day-ahead hay fever forecast for patients with grass-pollen-induced allergic rhinitis. *International Journal of*

- Biometeorology 2014; 58: 1047-1055.
- Dhiab AB, Mimoun MB, Oteros J, Garcia-Mozo H, Domínguez-Vilches E, Galán C, et al. Modeling olive-crop forecasting in Tunisia. *Theoretical and Applied Climatology* 2017; 128: 541-549.
- Durham OC. The volumetric incidence of atmospheric allergens: IV. A proposed standard method of gravity sampling, counting, and volumetric interpolation of results. *Journal of Allergy* 1946; 17: 79-86.
- Fernandez-Rodriguez S, Skjoth CA, Tormo-Molina R, Brandao R, Caeiro E, Silva-Palacios I, et al. Identification of potential sources of airborne Olea pollen in the Southwest Iberian Peninsula. *International Journal of Biometeorology* 2014; 58: 337-48.
- Galán C, Alcázar P, Oteros J, Garcia-Mozo H, Aira MJ, Belmonte J, et al. Airborne pollen trends in the Iberian Peninsula. *Science of the Total Environment* 2016; 550: 53-9.
- Galán C, Antunes C, Brandao R, Torres C, Garcia-Mozo H, Caeiro E, et al. Airborne olive pollen counts are not representative of exposure to the major olive allergen Ole e 1. *Allergy* 2013; 68: 809-12.
- Galán C, Cariñanos P, Alcázar P, Domínguez-Vilches E. Spanish Aerobiology Network (REA): management and quality manual: Servicio de Publicaciones, Universidad de Córdoba, 2007.
- Galán C, Domínguez-Vilches E. The capture media in aerobiological sampling. *Aerobiologia* 1997; 13: 155-160.
- Galán C, Smith M, Thibaudon M, Frenguelli G, Oteros J, Gehrig R, et al. Pollen monitoring: minimum requirements and reproducibility of analysis. *Aerobiologia* 2014; 30: 385-395.
- Gharbi D, Brighetti MA, Travaglini A, Trigo MM. Comparison between the counting methods used by two aerobiology networks in southern Europe (Spain and Italy). *Aerobiologia* 2017; 33: 87-92.
- Haberle SG, Bowman DM, Newnham RM, Johnston FH, Beggs PJ, Buters J, et al. The macroecology of airborne pollen in Australian and New Zealand urban areas. *PLoS One* 2014; 9: e97925.
- Hamilton JM, Tol RSJ. The impact of climate change on tourism in Germany, the UK and Ireland: a simulation study. *Regional Environmental Change* 2007; 7: 161-172.
- Hirst JM. An Automatic Volumetric Spore Trap. *Annals of Applied Biology* 1952; 39: 257-265.
- Hofmann F, Otto M, Wosniok W. Maize pollen deposition in relation to distance from the nearest pollen source under common cultivation-results of 10 years of monitoring (2001 to 2010). *Environmental Sciences Europe* 2014; 26: 24.
- Jochner-Oette S, Simmons M, Jetschni J, Menzel A. Impacts of land clearance by fire on spatial variation of mountain cedar pollen concentrations in Texas. *Landscape and Urban Planning* 2017; 162: 178-186.
- Kawashima S, Thibaudon M, Matsuda S, Fujita T, Lemonis N, Clot B, et al. Automated pollen monitoring system using laser optics for observing seasonal changes in the concentration of total airborne pollen. *Aerobiologia* 2017; 33: 351-362.
- Levetin E, Rogers CA, Hall SA. Comparison of pollen sampling with a Burkard Spore Trap and a Tauber Trap in a warm temperate climate. *Grana* 2000; 39: 294-302.
- Mandrioli P, Comtois P, Levizzani V. *Methods in aerobiology*: Pitagora Editrice Bologna, 1998.
- Maya-Manzano JM, Fernández-Rodríguez S, Silva-Palacios I, Gonzalo-Garijo A, Tormo-Molina R. Comparison between two adhesives (silicone and petroleum jelly) in Hirst pollen traps in a controlled environment. *Grana* 2017; 57: 137-143.
- Nakamori Y, Sawaragi Y. Interactive design of urban level air quality monitoring network. *Atmospheric Environment (1967)* 1984; 18: 793-799.
- O'Connor DJ, Healy DA, Hellebust S, Buters JT, Sodeau JR. Using the WIBS-4 (Waveband Integrated Bioaerosol Sensor) technique for the on-line detection of pollen grains. *Aerosol Science and Technology* 2014; 48: 341-349.
- Ogden EC, Raynor GS. A new sampler for airborne pollen: the rotoslide. *Journal of Allergy* 1967; 40: 1-11.
- Orlandi F, Oteros J, Aguilera F, Ben Dhiab A, Msallem M, Fornaciari M. Design of a downscaling method to

- estimate continuous data from discrete pollen monitoring in Tunisia. *Environmental Science: Process & Impacts* 2014; 16: 1716-25.
- Oteros J, Bartusel E, Alessandrini F, Núñez A, Moreno D, Behrendt H, et al. Artemisia pollen is the main vector for airborne endotoxin. *Journal of Allergy and Clinical Immunology* 2019; 143: 369-377.
- Oteros J, Buters J, Laven G, Roseler S, Wachter R, Schmidt-Weber C, et al. Errors in determining the flow rate of Hirst-type pollen traps. *Aerobiologia* 2017; 33: 201-210.
- Oteros J, Galan C, Alcazar P, Dominguez-Vilches E. Quality control in bio-monitoring networks, Spanish Aerobiology Network. *Science of the Total Environment* 2013a; 443: 559-565.
- Oteros J, Garcia-Mozo H, Alcazar P, Belmonte J, Bermejo D, Boi M, et al. A new method for determining the sources of airborne particles. *Journal of Environmental Management* 2015a; 155: 212-218.
- Oteros J, Garcia-Mozo H, Hervás-Martínez C, Galan C. Year clustering analysis for modelling olive flowering phenology. *International Journal of Biometeorology* 2013b; 57: 545-555.
- Oteros J, Orlandi F, Garcia-Mozo H, Aguilera F, Ben Dhiab A, Bonofiglio T, et al. Better prediction of Mediterranean olive production using pollen-based models. *Agronomy for Sustainable Development* 2014; 34: 685-694.
- Oteros J, Pusch G, Weichenmeier I, Heimann U, Moller R, Roseler S, et al. Automatic and Online Pollen Monitoring. *International Archives of Allergy and Immunology* 2015b; 167: 158-66.
- Portnoy J, Barnes C, Barnes CS. The National Allergy Bureau: pollen and spore reporting today. *Journal of Allergy and Clinical Immunology* 2004; 114: 1235-8.
- Rodríguez-Rajo FJ, Jato V, Fernández-González M, Aira MJ. The use of aerobiological methods for forecasting Botrytis spore concentrations in a vineyard. *Grana* 2010; 49: 56-65.
- Rojo J, Oteros J, Pérez-Badía R, Cervigón P, Ferencova Z, Gutiérrez-Bustillo AM, et al. Near-ground effect of height on pollen exposure. *Environmental Research* 2019a; 174: 160-169.
- Rojo J, Perez-Badía R. Effects of topography and crown-exposure on olive tree phenology. *Trees-Structure and Function* 2014; 28: 449-459.
- Rojo J, Picornell A, Oteros J. AeRobiology: the computational tool for biological data in the air. *Methods in Ecology and Evolution* 2019b; doi: 10.1111/2041 - 210X.13203.
- Saar, Meltsov. Passports of sampling sites in routine aerobiological monitoring of outdoor air. In: Clot B., Comtois P., B. E-G, editors. *Aerobiological Monographs, Towards a comprehensive vision*. 1. MeteoSwiss (CH) and University of Montreal (CA), Montreal, Canada, 2011, pp. 3.
- Šaulienė I, Šukienė L, Daunys G, Valiulis G, Vaitkevičius L, Matavulj P, et al. Automatic pollen recognition with the Rapid-E particle counter: the first-level procedure, experience and next steps. *Atmos. Meas. Tech. Discuss.* 2019; 2019; doi: 10.5194/amt-2018-432.
- Sikoparija B, Galán C, Smith M. Pollen-monitoring: between analyst proficiency testing. *Aerobiologia* 2017; 33: 191-199.
- Sikoparija B, Smith M, Skjoth CA, Radisic P, Milkovska S, Simic S, et al. The Pannonian plain as a source of Ambrosia pollen in the Balkans. *Int J Biometeorol* 2009; 53: 263-72.
- Simoleit A, Wachter R, Gauger U, Werchan M, Werchan B, Zuberbier T, et al. Pollen season of European beech (*Fagus sylvatica* L.) and temperature trends at two German monitoring sites over a more than 30-year period. *Aerobiologia* 2016; 32: 489-497.
- Smith M, Jager S, Berger U, Sikoparija B, Hallsdottir M, Sauliene I, et al. Geographic and temporal variations in pollen exposure across Europe. *Allergy* 2014; 69: 913-23.
- Smith M, Oteros J, Schmidt-Weber C, Buters J. An abbreviated method for the quality control of pollen counters. *Grana* 2019; 58: 185-190.
- Sofiev M, Ritenberga O, Albertini R, Arteta J, Belmonte J, Bonini M, et al. Multi-model ensemble simulations of olive pollen distribution in Europe in 2014. *Atmospheric Chemistry and Physics Discussions* 2017; 17: 12341-12360.
- Sofiev M, Siljamo P, Ranta H, Linkosalo T, Jaeger S, Rasmussen A, et al. A numerical model of birch pollen

- emission and dispersion in the atmosphere. Description of the emission module. *International Journal of Biometeorology* 2013; 57: 45-58.
- Sofiev M, Vira J, Kouznetsov R, Prank M, Soares J, Genikhovich E. Construction of the SILAM Eulerian atmospheric dispersion model based on the advection algorithm of Michael Galperin. *Geoscientific Model Development* 2015; 8: 3497-3522.
- Teranishi H, Kenda Y, Katoh T, Kasuya M, Oura E, Taira H. Possible role of climate change in the pollen scatter of Japanese cedar *Cryptomeria japonica* in Japan. *Climate Research* 2000; 14: 65-70.
- Thibaudon M, Šikoparija B, Oliver G, Smith M, Skjøth CA. Ragweed pollen source inventory for France– The second largest centre of *Ambrosia* in Europe. *Atmospheric Environment* 2014; 83: 62-71.
- VDI4252-4. Bioaerosole und biologische Agenzien – Ermittlung von Pollen und Sporen in der Außenluft unter Verwendung einer volumetrischen Methode für ein Messnetz zu allergologischen Zwecken. VDI-Richtlinie 4252 Blatt 4, Entwurf. VDI/DIN-Handbuch Reinhaltung der Luft, Band 1a. VDI/DIN, Beuth, Berlin, 2016.
- Ward Jr JH, Hook ME. Application of an hierarchical grouping procedure to a problem of grouping profiles. *Educational and Psychological Measurement* 1963; 23: 69-81.
- Werchan B, Werchan M, Mucke HG, Gauger U, Simoleit A, Zuberbier T, et al. Spatial distribution of allergenic pollen through a large metropolitan area. *Environmental Monitoring and Assessment* 2017; 189: 169.
- Zhang Y, Bielory L, Mi Z, Cai T, Robock A, Georgopoulos P. Allergenic pollen season variations in the past two decades under changing climate in the United States. *Glob Chang Biol* 2015; 21: 1581-1589.
- Ziello C, Böck A, Estrella N, Ankerst D, Menzel A. First flowering of wind - pollinated species with the greatest phenological advances in Europe. *Ecography* 2012a; 35: 1017-1023.
- Ziello C, Sparks TH, Estrella N, Belmonte J, Bergmann KC, Bucher E, et al. Changes to airborne pollen counts across Europe. *PLoS One* 2012b; 7: e34076.
- Zink K, Vogel H, Vogel B, Magyar D, Kottmeier C. Modeling the dispersion of *Ambrosia artemisiifolia* L. pollen with the model system COSMO-ART. *Int J Biometeorol* 2012; 56: 669-680.

Web references:

<https://www.zaum-online.de/pollen/pollen-monitoring-map-of-the-world.html>, 01-01-2019

<http://pollensense.com/>, 01-01-2019

<http://silam.fmi.fi>, 01-01-2019

www.swisens.ch, 01-01-2019

<http://www.schneefernerhaus.de/startseite.html>, 01-01-2019

<https://bwi.info/>, 05-05-2019

Supplementary Material

- Supplementary Material Figure S1.
- Supplementary Material Figure S2.
- Supplementary Material Figure S3.
- Supplementary Material Figure S4.
- Supporting data are freely available by petition. Please contact the corresponding author.

Figure captions

Figure 1. Footprint of System for Integrated modeLLing of Atmospheric coMposition (SILAM, <http://silam.fmi.fi>) for the birch pollen season of 2015 for the 27 ePIN locations.

Figure 2. A) Boxplot showing the Seasonal Pollen Integral (SPI) of the considered 13 pollen types at the 26 ePIN locations (all excluding DEERLA). **B)** Boxplots showing the start- and end date of the pollen season in 2015 (80% of annual pollen) in Bavaria.

Figure 3. ePIN test pollen network implemented during 2015 in Bavaria. This consisted of Hirst-type pollen traps at 27 locations under homogeneous monitoring conditions.

Figure 4. 4a) Dendrogram deriving from hierarchical clustering by use of Ward's method of Euclidean distances. **4b)** Network plot representing the position of the 26 considered ePIN stations (nodes) at a bi-dimensional space based on the 5 shortest Euclidean distances of each station (edges).

Figure 5. Final selected 8 locations for the ePIN network for permanent automatic pollen monitoring in Bavaria.

Supplementary figures:

Figure S1. Topographical map of Germany and Bavaria (CGIAR SRTM). **a)** Bavaria in the south of Germany. **b)** The 7 administrative districts of Bavaria (Regierungsbezirke).

Figure S2. a) Population density over Bavaria (CIESIN, 2010). **b)** Forest distribution over Bavaria (CORINE Land Cover, 2006). **c)** Average annual temperature (°C) (WorldClim, 1960-1990, Hijmans et al., 2005). **d)** Annual precipitation (mm/m²) (WorldClim, 1960-1990, Hijmans et al., 2005).

Figure S3. Equipment used during the building of ePIN testing network. **a)** a 1.5 m above ground level standard self-designed tower for supporting Hirst-type pollen trap. **b)** Central pollen lab at ZAUM (Munich), capable to manage 25 pollen stations / week. **c)** Draft of pollen slide, each slide corresponds to one independent day. **d)** 12.5 mm net micrometer used for limiting the sub-sampling to 9% of the slide by analysing 4 lines, a net micrometer of 10 mm was also allowed following the national German standards (4 lines = 7% subsampling) **e)** Examples of own developed equipment: flowmeter without air resistance (Oteros et al., 2017), ruler with support for cutting samples, vaseline coating machine with a rotating motor, guide for drawing standard limits into the slides. **f)** Computer program enabling keyboards for pollen counting. Screenshot of the online Aerex-ZAUM database to store the counting data.

Figure S4. Elbow plot suggesting the appropriate number of clusters of the dataset.

Table 1. Selected locations for ePIN. Population coverage (%) within 30-km from each station is given.

Code	Site Location (German)	Latitude (°)	Longitude (°)	Altitude (m a.s.l.)	Population coverage (%)
DEALTO	Altötting	48.23	12.68	398	2.5
DEAUGS	Augsburg	48.33	10.90	497	6.4
DEBAMB	Bamberg	49.90	10.89	238	3.6
DEBAYR	Bayreuth	49.94	11.53	419	2.8
DEBERC	Berchtesgaden	47.64	13.01	573	0.8
DEBIED	München	48.16	11.59	510	15.2
DEDONA	Donaustauf	49.04	12.21	425	3.4
DEERLA	Erlangen	49.60	11.01	284	9.7
DEFEUC	Feucht (Nürnberg)	49.38	11.20	365	8.9
DEGAIS	Gaissach	47.75	11.58	717	2.7
DEGARM	Garmisch-Partenkirchen	47.49	11.10	821	1.2
DEHOF	Hof	50.32	11.90	531	1.9
DEKITZ	Kitzingen	49.74	10.14	246	3.6
DEKOES	Kösching	48.82	11.51	391	3.2
DELANDS	Landshut	48.54	12.14	397	3.0
DEMARK	Marktheidenfeld	49.85	9.63	216	3.4
DEMIND	Mindelheim	48.04	10.50	610	3.3
DEMUNC	München	48.13	11.56	538	15.1
DEMUST	Münnerstadt	50.25	10.18	347	2.6
DEOBER	Oberjoch	47.52	10.40	870	1.7
DEOETT	Oettingen	48.96	10.60	431	1.8
DEPASS	Passau	48.56	13.44	318	2.1
DETROS	Trostberg	48.03	12.56	483	2.7
DEUFS	Umwelt Forschungsstation Schneefernerhaus (UFS)	47.42	10.99	2650	0.8
DEVIEC	Viechtach	49.08	12.87	459	2.0
DEWEID	Weiden	49.68	12.17	403	2.0
DEZUSM	Zusmarshausen	48.40	10.61	483	5.7


Table 2. Standard conditions for homogeneous pollen monitoring in the network.

CONDITIONS FOR POLLEN MONITORING		
Logistic	Trap location	Emission sources
Safety location	Flat and horizontal surface	Absence of overrepresentation of some species in surrounding 500 m area
Easy access	Higher than surrounding roofs and other wind walls	Absence of anemophilous sources in surrounding: uncut grass areas (no in 50 m) and birch/olive trees (no in 100 m).
Access to electric network	Not at ground level. Between 9 m and 15 m from the ground (on a roof or an elevation tower). This criterion aims to build the whole Bavarian network under homogeneous conditions to have a more uniform information over a greater area in comparison to measurements on the ground with an extreme influence of plants nearby.	Absence of proximity to non-biological and biological particle sources of high emission (e.g. waste plant)
Access to internet (For 3G systems)	Not placed at the edge of a building (> than 2 meters) to avoid turbulent flow	Absence of proximity to wind distortion sources (e.g. solar panels, refrigeration systems...)
Temporal sustainability	Elevated more than 150 cm from the roof or elevation surface	Consider whether land use change will have an effect on pollen concentrations in future

Table 3. Sub-clusters and selection criteria. Within each subcluster, the station with the highest population was selected. In the case of draw, the selection criteria were applied for tiebreaking and selecting the most representative station within the sub-cluster.





Ward Sub-Cluster	Code	% of covered population	Selected	Criteria
1.1	DEAUGS	6.4	No	Standard Criteria - <70km apart from the closest station (DEMUNC)
1.1	DEMIND	3.3	Yes	Standard Criteria
1.2	DETROS	2.7	No	Population draw
1.2	DEALTO	2.5	Yes	Population draw, Closer to the border
1.2	DEPASS	2.1	No	Standard Criteria
1.3	DEBIED	15.2	No	Population draw
1.3	DEMUNC	15.1	Yes	Population draw, Closer to the border
1.3	DEDONA	3.4	No	Standard Criteria
1.3	DEKOES	3.2	No	Standard Criteria
1.3	DEOETT	1.8	No	Standard Criteria
1.4	DEFEUC	8.9	Yes	Standard Criteria
1.4	DEWEID	2	No	Standard Criteria
2.1	DEVIEC	2	Yes	Standard Criteria
2.1	DEBERC	0.8	No	Standard Criteria
2.2	DEGAIS	2.7	No	Standard Criteria - <70km apart from the closest station (DEMUNC)
2.2	DEOBER	1.7	No	DEOBER was already selected for manual monitoring in a parallel manual network
2.2	DEGARM	1.2	Yes	Standard Criteria
2.2	DEUFS	0.8	No	Standard Criteria
3	DEZUSM	5.7	No	Outlier
3	DELAND	3	No	Outlier
3	DEMUST	2.6	No	Outlier
4.1	DEBAMB	3.6	No	Population draw
4.1	DEKITZ	3.6	No	Population draw
4.1	DEMARK	3.4	Yes	Population draw, Closer to the border
4.2	DEBAYR	2.8	No	Standard Criteria - <70km apart from the closest station (DEFEUC)
4.2	DEHOF	1.9	Yes	Standard Criteria
---	DEERLA	9.7	No	Standard Criteria


Footprint:sum



ePIN testing network (2015), based on manual-Hirst pollen trap (PID: Stiftung Deutscher Polleninformationsdienst)




-  abandoned PID station, temporarily reactivated for ePIN test-network
-  PID-station currently in service
-  other, non-PID operated station
-  ePIN test-station 2015




0 55 110 220
Kilometer



a) Ward cluster rescaled distance







Reordered dendrogram from
hclust(d = norm.ch, method = "ward.D")

b) Euclidean distances network











ePIN network

 Electronic Pollen Monitor (ePOMO)



0 55 110 220 Kilometers



Test Network

Clustering

ePIN Network


a)




0 25 50 100 150 Kilometers







b)



0 15 30 60 90 Kilometers





f)

Key	in	'ON'	'OFF'	'ON'	'OFF'	Pollen	Pollen Counts
Key	Key	Key	Key	Key	Key		
A	Enter/Rt	1	F1	1		<i>Betula</i>	
S	L	2	F2	2		<i>Taxus</i>	
D	K	3	F3	3		<i>Fraxinus</i>	
F	J	4	F4	4		<i>Alnus</i>	
G	H	5	F5	5		<i>Corylus</i>	
V	-	6	F6	6		<i>Populus</i>	
R	.	7	F7	7		<i>Urtica</i>	
E	,	8	F8	8		<i>Picea</i>	
R	M	9	F9	9		<i>Fagus</i>	
T	N	0	F0	0		<i>Pinus</i>	
Z	Y	UP	RT	Y		<i>Piceae</i>	
X	U	DOWN	F12	U		<i>Tilia</i>	
C	I	LEFT	F13	I		<i>Artemisia</i>	
V	O	RIGHT	F14	O		<i>Androsida</i>	
B	P	Enter (Num)	F16	P		<i>Indeterminate (the rest of</i>	

0:00h-12:00h
Line 1

0:00h-12:00h
Line 2

0:00h-12:00h
Line 3

0:00h-12:00h
Line 4

12:00h-24:00h
Line 1

12:00h-24:00h
Line 2

12:00h-24:00h
Line 3

12:00h-24:00h
Line 4

INFORMATION	
Responsible for Creating	Jesse Dierckx
Monitoring Station	DEZUSIM
Date	2015-05-15
Year since launch or use	
Microsites	
at (x400 magnification), i.e. usually between 0.45mm and 0.65mm.	0.243

Aeroallergen Exposome Database

Overview Import Logout

Editor Export

Filter

Project: ePIN (2015-01-01 - 2015-12-31)

Locations: DEALTO, DEAU2, DEAU6S, DEBAMB, DEBAYR, DEBERC, DEBIE2, DEBIE2 (Hirst), DEBIE2-Lohstr

Date Range: From: To:

There are no available values for the selected locations.